

03/27/98
JCS64 U.S. PTO

A
Dkt. 47765-C/JPW/EMW

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Collin J. Weber et al.
Appln No. : Not Yet Known
Filing Date : Herewith
For : METHOD OF INHIBITING IMMUNE SYSTEM DESTRUCTION
OF TRANSPLANTED VIABLE CELLS

1185 Avenue of the Americas
New York, New York 10036
March 27, 1998

Assistant Commissioner for Patents
Washington, D.C. 20231

Box: Patent Application

EXPRESS MAIL CERTIFICATE OF MAILING
FOR ABOVE-IDENTIFIED APPLICATION

"Express Mail" mailing label number: EH 854 722 048 US

Date of Deposit: March 27, 1998

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Printed Name: John P. White

Respectfully submitted,

Albert Wai Kit Chan

John P. White
Registration No. 28,678
Albert Wai-Kit Chan
Registration No. 36,479
Attorneys for Applicants
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

March 27, 1998

Honorable Assistant Commissioner for Patents
 Washington, D.C. 20231
 Re: Patent Application
 of R:

Transmitted herewith for filing are the specification and claims of the patent application of:

Collin J. Weber, Mary K. Hagler, Peter S. Linsley, Judith A. Kapp, Susan A. Safley for Inventor(s)

METHOD OF INHIBITING IMMUNE SYSTEM DESTRUCTION OF TRANSPLANTED VIABLE CELLS
 Title of Invention

Also enclosed are:

☒ 56 sheet(s) of ☐ informal ☒ formal drawings.

☐ Oath or declaration of Applicant(s).

☐ A power of attorney

☐ An assignment of the invention to _____

☒ A Preliminary Amendment

☐ A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED		NUMBER EXTRA*		RATE		FEE	
					SMALL ENTITY	OTHER ENTITY	SMALL ENTITY	OTHER ENTITY
Total Claims	34 -20	=	14	X	\$11.00	\$22.00	= \$	\$308
Independent Claims	2 -3	=	0	X	\$41.00	\$82.00	= \$	\$ 0
Multiple Dependent Claims Presented: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No					\$135.00	\$270.00	= \$	\$ 0
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE		\$ 395	\$ 790
					TOTAL FEE		\$	\$1098

Applicants: Collin J. Weber et al.
U.S. Serial No.: Not Yet Known
Filed: Herewith

Letter of Transmittal
Page 2

X A check in the amount of \$_____ to cover the filing fee.

_____ Please charge Deposit Account No. _____ in the amount of \$_____.

X The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:

X Filing fees under 37 C.F.R. \$1.16.

X Patent application processing fees under 37 C.F.R. \$1.17.

_____ The issue fee set in 37 C.F.R. \$1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. \$1.311(b).

X Three copies of this sheet are enclosed.

A certified copy of previously filed foreign application No. _____
_____ filed in _____ on _____.

Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.

X Other (identify) Express Mail Certificate of Mailing bearing label No.
EH 854 722 048 US dated March 27, 1998 and a loose set of
figures.

Respectfully submitted,

Albert Wai Kit Chan

John P. White
Registration No. 28,678
Albert Wai-Kit Chan
Registration No. 36,479
Attorneys for Applicants
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

By this Preliminary Amendment, claims 24-42 have been canceled without prejudice to applicants' right to pursue the subject matter


Applicants: Collin J. Weber, et al.
U.S. Serial No. : Not Yet Known
U.S. Filing Date: Herewith
Page 2

of these claims in the future. Accordingly, only claims 1-23 and 43-53 are now pending.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicants' undersigned attorney invites the Examiner to telephone at the number provided below.

No fee is deemed necessary in connection with the filing of this Preliminary Amendment. However, if any fee is required, authorization is hereby given to charge the amount of such fee to Deposit Account No. 03-3125.

Respectfully submitted,



John P. White
Registration No. 28,678
Albert Wai-Kit Chan
Registration No. 36,479
Attorneys for Applicants
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

1344260-9984060

**Application
for
United States Letters Patent**

To all whom it may concern:

Be it known that We,
Collin J. Weber, Mary K. Hagler, Peter S. Linsley, Judith A. Kapp and Susan A. Safley
have invented certain new and useful improvements in

METHOD OF INHIBITING IMMUNE SYSTEM DESTRUCTION OF TRANSPLANTED VIABLE CELLS

of which the following is a full, clear and exact description.

METHOD OF INHIBITING IMMUNE SYSTEM DESTRUCTION OF TRANSPLANTED VIABLE CELLS

This application is a continuation-in-part of PCT International Application No. PCT/US96/15577, filed September 27, 1996 which claims the benefit of U.S. 5 Provisional Application No. 60/004,375, filed September 27, 1995, the contents of which are hereby incorporated by reference.

The invention disclosed herein was made with Government
10 support under NIH Grant Nos. RO1-DK39088 and RO1DK53057.
Accordingly, the U.S. Government has certain rights in this
invention.

Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be found at the end of this application, preceding the claims.

Background of the Invention

There is a critical need for better insulin replacement therapy to circumvent the complications of insulin-dependent diabetes mellitus (IDDM). Our goal is to develop techniques for transplantation of microencapsulated, xenogeneic islets to provide a durable, physiological source of insulin to diabetic patients. It has previously been shown that microcapsules are biocompatible and that xenogeneic islet grafts contained in microcapsules functioned indefinitely in the peritoneal cavity of mice with streptozotocin-induced (SZN) diabetes. Thus, microcapsules may be intact and stable *in vivo* and factors

that may be required for long-term survival and function of the xenogeneic islets are accessible. The microcapsules serve as a mechanical barrier that prevents cell-to-cell contact between recipient lymphocytes and donor islets.

5 The mechanical barrier primarily prevents host sensitization rather than protecting the graft from immune destruction, because encapsulated islets are very rapidly destroyed by recipients that are presensitized to the islet donor cell antigens. Similarly, encapsulated xenogeneic

10 islets were rejected (in two weeks) by NOD mice, which is possibly due to presensitization of NODs to islet antigens. Xenografts undergoing rejection in NOD mice were surrounded by large numbers of activated macrophages and immunoglobulins, with IL-1 α , TNF α , both documented by

15 immunocytochemistry, and IL-4 messenger RNA detected by RT-PCR. We postulate that NOD rejection is initiated by donor antigens that are secreted by or shed from the encapsulated islets and which are processed via the MHC (major histocompatibility complex) class II pathway by host APC

20 (antigen presenting cells). These APC activate NOD CD4⁺ T cells that develop into a Th2 response, with donor islet destruction occurring via cytokine-mediated events.

We have also been able to improve the microencapsulation

25 process to permit long-term survival of concordant, rat islet xenografts, even in NOD mice. Furthermore, we have found that blockade of NOD co-stimulatory molecules with CTLA4Ig significantly prolongs survival of discordant, rabbit islet xenografts for up to 200 days. Thus, we have

30 been able to overcome problems associated with transplanting encapsulated islet xenografts into autoimmune diabetic recipients.

Insulin-Dependent Diabetes Mellitus

35 The last several years have witnessed a remarkable increase

2025 RELEASE UNDER E.O. 14176

in or knowledge of the effects of therapies for insulin-dependent diabetes mellitus (IDDM). The Diabetes Control and Complications Trial (DCCT) found that intensive insulin therapy delayed the onset and slowed progression of
5 retinopathy, nephropathy, and neuropathy in patients with IDDM (1). Unfortunately, intensive insulin therapy is not appropriate for many IDDM patients; and even with careful monitoring, DCCT patients had increased episodes of severe hypoglycemia (1). Ironically, results of the DCCT support
10 the rationale for pancreas and islet transplantation. Since the inception of islet transplant experiments, it has been the hope that such grafts might supply insulin more homeostatically than exogenous insulin can, and that 'near-normal' modulation of carbohydrate metabolism might prevent
15 the secondary complications of IDDM (2). Clinical pancreas allografts have improved outcomes with the advent of combination immunosuppression; and near normal of glucose homeostasis follows most pancreatic allo- and auto-grafts (3). However, the first-year mortality of a human
20 pancreatic allograft remains high (10%), immunosuppression is required, and only limited numbers of clinical whole-organ pancreatic transplants are being done worldwide (2,4,5).

25 **The Rationale for Microencapsulated Islet Xenografts**

Islet transplantation is an attractive therapy for patients with IDDM, since problems related to the exocrine pancreas may be avoided. However, allografts of donor human islets have not been successful long-term (3); and availability
30 and yield of human islets are limited. Therapeutic islet transplants for large number of patients almost certainly will require donor islets harvested from animals (xenografts) (2,4).

35 The optimal source of xenogeneic islets for clinical use

remains controversial. Islets have been isolated from subhuman primates and xenografted into immunosuppressed, diabetic rodents, with short-term reversal of diabetes (6). However, there are significant ethical issues surrounding use of primates, Other promising sources are porcine, bovine, canine, and rabbit islets, which function remarkably well, (i.e., maintaining normoglycemia) in diabetic rodents until transplant rejection occurs (7-11). Long-term human, bovine and porcine islet xenograft survival has been documented in nude mice and rats, suggesting that sufficient islet-specific growth factors are present in xenogeneic recipients (2,12-17). For sociologic/ethical reasons, canine islets are not clinically appropriate. Porcine islets are both difficult to isolate (intact) and to maintain *in vitro*; nevertheless, they are extremely promising for eventual clinical application (18-21). Isolation of bovine islets is technically easier (than porcine islets), and calf islets are glucose-responsive (22). Recently, large scale rabbit islets isolation has been developed (23) (see Preliminary Studies). Rabbit pancreas is an attractive source of islets. Rabbit, like porcine insulin, differs from human insulin at only one amino acid, and rabbit islets are glucose responsive (22,24). In addition, most humans do not possess natural anti-rabbit antibodies, which might improve the possibility of preventing xenograft rejection (25). It is currently feasible to consider isolation of 1,000,000 donor islets/per human diabetic recipient from either calves, pigs or rabbits, utilizing multiple donors.

The most significant obstacle to islet xenotransplantation on human IDDM is the lack of an effective immunosuppressive regiment to prevent cross-species graft rejection (2,26-28). Recently, it has been reported that human islets will survive long-term in SZN-diabetic mice treated either with

anti-CD4 antibody (16) or CTLA4Ig (a high affinity fusion protein which blocks CD28-B7 interactions) (12), or by exposure of donor islets to purified high affinity anti-HLA (ab)₂ (29). However, with the exception of these studies, indefinite survival of islet xenografts has rarely been achieved, except with the aid of porous, mechanical barriers. Both intra- and extra-vascular devices are under development. However, potential clinical complications, such as bleeding, coagulation, and bioincompatibility mitigate against their current use in diabetic patients (30,31). For example, acrylic-copolymer hollow fibers placed subcutaneously maintained viability of human islet allografts for two weeks (50 islets per 1.5 cm fiber) (65,000 M.W. permeability) (32).

However, to implant 500,000 islet would require >150 meters of these hollow fibers, which is not clinically feasible.

One of the most promising islet envelopment methods is the polyamino acid-alginate microcapsule. A large number of recent studies have shown that intraperitoneal xenografts of encapsulated rat, dog, pig or human islets into streptozotocin-diabetic mice or rats promptly normalized blood glucose for 10-100+ days (7,19,33-39). Long-term normalization of hyperglycemia by microencapsulated canine islet allografts, porcine islet xenografts, and one human islet allograft has been reported (21,40-42). The mechanisms by which microcapsules protect islet xenografts from host destruction are not fully understood. However, it has been suggested that prohibition of cell-cell contact with host immunocytes is important (30,35). The marked prolongation of widely unrelated encapsulated islet xenografts in rodents with induced diabetes has prompted studies in animals with spontaneous diabetes.

Nonobese diabetic (NOD) mice develop diabetes spontaneously, beginning at approximately twelve weeks of age. NOD mice are the most appropriate model for studying the feasibility of islet xenotransplants because their disease resembles human IDDM in several ways. Macrophage, dendritic cell and lymphocytic infiltration of islets can be detected as early as four weeks of age and precedes overt hyperglycemia (43-46). NOD diabetes is T lymphocyte-dependent (43-45); and it is associated with (MHC) Class II genes (47-50). Cytotoxic T cells and antibodies specific for beta cells or for insulin have been identified, characterized and cloned from NOD mice (44,45,51-55). Loss of tolerance to islet antigens in NODs correlates with appearance of Th1 immune responses to glutamic acid decarboxylase, a factor which has been reported to be a primary auto-antigen in human IDDM (5,657). The disease can be induced in non-diabetic, syngeneic mice by transfer of both CD8⁺ and CD4⁺ T cells or T-cell clones from diabetic NODs (44,52,55,58); and inhibition of NOD macrophages or CD4⁺ T lymphocytes or treatment with anti-Class II monoclonal antibodies prevents or delays diabetes onset in NOD mice (59,50). Defects in NOD macrophages, C5 complement and NK cell function have been reported (61). It has been suggested that helper T-cells function to activate CD8⁺ cells, which damage beta cells by direct cytotoxic attack. However, some recent studies have suggested that beta cell killing may be indirect, from a nonspecific inflammatory response which initially involves CD4⁺ cells, but also includes infiltrating macrophages, which release cytokines and oxygen free-radicals (particularly nitric oxide), known beta cell toxins (62-65). Because of similarities to IDDM, NOD mice are the best model in which to study islet xenografts.

Recently, the Scid mutation has been back-crossed onto the NOD background, resulting in immuno-deficient NOD-Scid mice (66-69). These mice homologous for the Scid mutation, which results in an inability to rearrange T-cell receptor and immunoglobulin genes (66,67). The consequence is an absence of T and B-lymphocytes. These mice do not develop diabetes spontaneously; but they may be rendered diabetic with multiple low-dose streptozotocin (MLD-SZN) regimens, making them an optimal model for adoptive transfer experiments (67-69). NOD-Scids express NOD MHC genes and other genes that are relevant for development of the disease. They mount robust macrophage and limited NK-cell responses, but are functionally T- and B-lymphocyte deficient (69).

15

Islet Xenografts into Diabetic NOD Mice

Unlike mice with SZN-induced diabetes, diabetic NOD mice rapidly reject unencapsulated islet xenografts, allografts and isografts (7,8,10,19,33,56,70,71). Conventional immunosuppressive regimens have little effect on this reaction (10,71-73). Treatment of NOD recipients with monoclonal antibodies directed against CD4⁺ helper T lymphocytes or FK506 prolongs islet graft function (from 5 to 25 days) (7,8,10,73); but long-term islet graft survival in NODs has not been reported.

Several laboratories have reported that intraperitoneal microencapsulated islets (allo- and xeno-geneic) function significantly longer than non-encapsulated controls, but eventually are destroyed also by recipients with spontaneous (autoimmune) diabetes (NOD mice or BB rats) (7,9,19,33,35,70,74-78). Rejection is accompanied by an intense cellular reaction, composed primarily of macrophages and lymphocytes, which entraps islet-containing microcapsules and recurrence of hyperglycemia within 21

days, in both NOD and BB recipients (7,19,74,76,77). The mechanism of encapsulated islet rejection by animals with spontaneous diabetes remains incompletely understood, but the fact that it rarely occurs in mice with induced (SZN) diabetes suggests that anti-islet autoimmunity may be involved in islet graft destruction.

Mechanisms of NOD Destruction of Encapsulated Islet Xenografts: Macrophages, T-Cells, and Cytokines

It has been suggested by several investigators that microcapsules, like other bioartificial membrane devices promote survival of xenogeneic and allogeneic islets by: (A) preventing or minimizing release of donor antigen(s), thereby reducing host sensitization, and/or (B) preventing or reducing host effector mechanisms (i.e. T-cell contact, anti-graft antibody binding, cytokine release).

Most studies of rejection of islets in microcapsules and other membrane devices have focused on effector mechanisms. For example, Halle (35) and Darquy and Reach (79) reported that microcapsules protected donor islets from host immunoglobulins, specifically human anti-islet antibodies and complement effects, *in vitro*. Although complement components, are too large ($>150,000$ Kd) to enter conventional poly-L-lysine microcapsules, it is possible that antibodies combine with shed donor antigens forming complexes which bind to FcR of macrophages *in vivo* (in the peritoneal cavity) which could initiate cytokine release causing encapsulated islet destruction (80). Complement could facilitate binding of complexes to macrophages via the C3b receptor or by the release of chemotactic peptides that could increase the number of macrophages.

Involvement of NOD T-lymphocytes in rejection of encapsulated islets has been proposed by Iwata, et al.

A prominence of macrophages/monocytes in peri-microcapsular infiltrates of encapsulated islet allografts and xenografts in NOD mice and BB rats has been reported (7,33,36,74,76-78,84). Cytokines known to be products of macrophages, including IL-1 and TNF (62,77,85,86), may be involved in destruction of encapsulated islets. Both IL-1 and TNF have been reported to reduce insulin secretion and cause progressive damage of islet cells *in vitro* (58,62-64,85-87). Cytokine-mediated injury might occur directly or indirectly, by activation of an intraperitoneal inflammatory response (30,77). Recently, it has been reported by Dr. J. Corbett (IPITA conf. 6/95), that there are as many as ten macrophages within each islet. IL-1 induces nitric oxide synthase (NOS) (63-65), with resultant generation of nitric oxide (NO), which causes injury to mitochondria and to DNA in beta cells (63-65). Furthermore, this pathway of islet damage is worsened by TNF (88,89). Theoretically, macrophages from within donor islets and host peritoneal cavity or within the down islets could be involved in cytokine-mediated damage to encapsulated islets.

[illegible][illegible]

060907

however, subsequent re-exposure of T-cells to peptide, mitogen, etc. will result in proliferation in the absence of costimulation. (95).

5 CTLA4 is a cell surface protein that is closely related to CD28; however, unlike CD28, CTLA4 is expressed only on activated T-cells. B7-1 has a high affinity for CLTA4 than CD28; and it has been suggested that CTLA4 may modulate functions of CD28 (97,103,104). CTLA4Ig is a recombinant
10 soluble fusion protein, combining the extracellular binding domain of the CTLA4 molecule with constant region of the IgG₁ gene. Both human and murine CTLA4Ig have been shown to inhibit T-lymphocyte responses in mice (141,142). Administration of CTLA4Ig to mice has been shown to induce
15 antigen-specific unresponsiveness (in a murine lupus model) (97,99,105) and long-term acceptance of murine cardiac allografts (106,107). In addition, Lenschow, et al., found that it induced tolerance to human islets in SZN-diabetic mice (12). CTLA4Ig has also been reported to
20 reduce the incidence of diabetes in NODs (108). There are no reports of effects of CTLA4Ig on islet graft survival in spontaneously-diabetic recipients, such as NOD mice. However, our studies show that CTLA4Ig significantly prolongs survival of encapsulated rabbit islets in NOD
25 recipients.

Recent studies have further illuminated helper T-cell-APC interactions, with recognition of the importance of binding of the APC-CD40 antigen to its ligand, GP39, on helper T-
30 cells (109,110). A monoclonal hamster anti-murine GP39 antibody (MR1) blocks helper T-cell interactions with APCs, macrophages, effector T-cells and B-lymphocytes (109,110). Dr. A. Rossini has reported recently (IPITA conf. 6/95) that MR1 plus B7 negative donor spleen cells day 7 allows
35 long-term survival of both allo- and xeno-geneic islets in

SZN-diabetic mice.

The Immunogenicity of Encapsulated Islets and Mechanisms of Graft of Destruction

5 Empty microcapsules have been reported to elicit no cellular responses (33,35,36). On the other hand, others have found reactions to empty capsules, (30,76,77,111,112). Impurities in reagents such as contamination with endotoxin or high concentrations of mannuronate most likely
10 contribute to bioincompatibility (113). It is apparent that some formulations of poly-l-lysine microcapsules are biocompatible and some are not. Until standardized reagents are available, immunologic studies are microencapsulated islets can only be interpreted when
15 investigators include empty microcapsule controls which document their biocompatibility.

Recently, de Vos, et al. (114) reported incomplete encapsulation or actual protrusion of islets through
20 microcapsule membranes in some microcapsules, and suggested this biomechanical imperfection is one factor in microcapsule destruction. Similar observations have been made by Chang (115), who found incorporation of islets and hepatocytes within the walls of poly-l-lysine alginate
25 microcapsules. Several other investigators have published photomicrographs of encapsulated islets showing obvious entrapment of islets in capsules, walls, but did not comment on this problem (35,116,117). Incomplete encapsulation would be anticipated to result in premature
30 capsule fracture and exposure of donor islets to host cells; but there are no reports analyzing this as a source of donor antigen exposure, sensitization and host.

Relatively few studies have focused on the role of donor
35 islet antigen(s) released from microcapsules in initiating

55220" 59861060

host immune responses. Ricker, et al. (33) reported similar, intense cellular reactions by NOD mice to rat insulinoma, hepatoma and pheochromocytoma cell lines in microcapsules and concluded that the NOD immune reaction
5 was not islet-specific. Horcher, et al. (36) reported 15-week survival of 6/7 encapsulated Lewis rat islet isografts, compared to failure of 8/10 encapsulated Wistar-to-Lewis islet allografts within 56 days. Isograft biopsies showed viable islets, intact capsules and no
10 pericapsular immune reaction (36), while biopsies of failed allografts revealed pericapsular cellular responses and nonviable islets. This is the only report in the literature with encapsulated islet isograft controls. Although the Lewis rat model is not one with autoimmune
15 diabetes, the results are significant, and suggest that donor antigen(s) are the stimulus for subsequent host responses.

09049865-032798

Summary of the Invention

This invention provides a method of inhibiting viable cells transplanted into a subject from being destroyed by the subject's immune system which comprises: a) containing the
5 viable cells, or tissue comprising the viable cells, prior to transplantation within a device comprising a semipermeable membrane; and b) treating the subject with a substance which inhibits an immune-system costimulation event in an amount effective to inhibit the subject's
10 immune system from responding to said contained cells or tissue.

In one embodiment, the substance which inhibits an immune-system costimulation event is CTLA4. Accordingly, this
15 invention further provides a method of inhibiting viable cells transplanted into a subject from being destroyed by the subject's immune system which comprises: a) containing the viable cells, or tissue comprising the viable cells, prior to transplantation within a device comprising a
20 semipermeable membrane; and b) treating the subject with CTLA4 in an amount effective to inhibit the subject's immune system from responding to said contained cells or tissue.

25 This invention also provides a method of treating diabetes in a subject which comprises: a) containing viable insulin-producing cells, or tissue comprising viable insulin-producing cells, within a device comprising a semipermeable membrane so as to obtain contained viable insulin-producing
30 cells; b) transplanting contained viable insulin-producing cells obtained in step (a) into the subject in an amount effective to treat diabetes in the subject; and c) treating the subject with a substance which inhibits an immune-system costimulation event in an amount effective to
35 inhibit the subject's immune system from responding to an

004535-03298
PAGE 5

amount of contained viable insulin-producing cells
according to step (b).

0304935 032793
0307220 03084060

immunocytochemistry demonstrates 5-10% beta cells. Approx. 400X.

5 Figure 17: Neonatal islet in microcapsule, biopsied
 day # 103 from SZN-diabetic NOD-Scid mouse.
 anti-insulin immunohistochemistry, showing
 intensely insulin-positive beta cells,
 occupying approximately 80% of islet.
10 Approx. 400X. Arrow points to outer
 surface of microcapsule membrane.

15 Figure 18: Non-encapsulated intrasplenic/portal
 neonatal porcine islet xenograft in
 streptozotocin diabetic NOD-Scid mouse.
 Biopsies (not shown) revealed viable
 porcine islets in both liver and splenic
 parenchyma.
 N=1
 T=Transplant
20 S=Sacrificed for biopsies of spleen and
 liver

25 Figure 19: Intraperitoneal microencapsulated neonatal
 porcine islet xenograft into
 streptozotocin-diabetic NOD-Scid mouse.
 Biopsied day #103 (see Fig. 20).
 N=1
 T=Transplant
30 S=Sacrificed

35 Figure 20: Neonatal porcine islet in microcapsule,
 biopsied day #103 after xenotransplantation
 to SZN-diabetic NOD-Scid mouse. H & E, X
 400. Arrow points to inner surface of
 microcapsule membrane.

004930 027000

- 86280"59867060
- 5 Figure 21: Encapsuled Neonatal Porcine Islet
Xenografts (N=5) in NODs, treated with
CTLA4Ig, 200 μ g i.p. Q.O.D., x 20 days.
NOD 880 was biopsied at day #101 (see Fig.
22).
S=Sacrificed for biopsy
(---)=Graft failure
- 10 Figure 22: Microencapsulated neonatal porcine islet,
biopsied 101 days after xenotransplantation
i.p. to spontaneously diabetic NOD mouse.
CTLA4Ig, 200 μ g i.p. Q.O.D., days # 0-21.
Arrow points to inside of intact
microcapsule wall. No pericapsular NOD
cellular response. H. & E. x200.
- 15
- 20 Figure 23: Adjacent section of same biopsy Anti-
insulin immunocytochemistry demonstrates
that most cells are insulin-positive beta
cells. x400.
- 25 Figure 24: Intraperitoneal microencapsulated neonatal
porcine islet xenografts in NOD mice
treated with CTLA4Ig* , which does not fix
complement.
- 30 Figure 25A-D: Spleen cells were cultured at 2×10^6 cells/ml
in 96-well plates with no antigen, 10 empty
capsules, 10 capsules containing neonatal
pig islets, 4×10^3 neonatal pig islets that
were unirradiated or irradiated with 2000R.
Spleen cells were obtained from normal NOD
mice (panel A); diabetic NOD mice (panel
B); diabetice NOD mice that were
35 transplanted with encapsulated, neonatal

pig islets and injected with CTLA4Ig (panel C) or mutant CTLA4Ig* (panel D) as described in Fig. 24. After 48 hrs incubation, ³H-TdR was added and the cells harvested 18 hrs later. Results represent the average \pm SD of triplicate cultures.

Figure 26:

Lymphokine production in cultures of spleen cells from the mice described in Fig. 24 were determined by ELISA. Spleen cells from normal or diabetic NOD mice were cultured with unirradiated neonatal, pig islets as described in Fig. 24. Supernatant fluids were harvested after 24 hrs of incubation and assayed for IL-4, IL-10 and IFN γ using a sandwich ELISA and the appropriate recombinant cytokines as standards.

Figure 27:

Model of immune response to micro encapsulated, xenogeneic islets by autoimmune, NOD mice. Secreted insulin clearly crosses the membrane of double walled microcapsules and regulated glucose levels in engrafted mice. 1): Potentially, other donor proteins or protein fragments of less than 100,000mw (AgX) that are shed or secreted by islets diffuse out of microcapsules and are endocytosed by dendritic cells. 2): Dendritic cells process proteins via the MHC class II pathway and present peptide X complexed with class II and co-stimulatory molecules to CD4⁺ T cells. In the presence of the appropriate cytokines, CD4⁺ T cells are

activated and develop into Th2 cells that express CD40L (GP39). B cells with surface IgM that bind AgX endocytose and process it into peptides that bind MHC class II which are expressed on the surface of B cells. Th2 specific peptide X complexed with class II binds B cells and the interaction of CD40 with CD40L (GP39) causes the activation of B cells. 3): Activated B cells mature into plasma cells under the direction of Th2 lymphokines. 4): Plasma cells secrete specific antibody that forms complexed with AgX. 5): Binding of complexes to FcR activated macrophages to secrete a variety of mediators including IL-1, TNF α and nitric oxide (NO), all of which have toxic effects on islets and all of which are small enough to cross the double-walled microcapsules.

Figure 28A-B: (A) Neonatal islet in microcapsule, biopsied day 103 from peritoneal cavity of SZN-diabetic NOD-Scid mouse (H&E) (approx. 240X). Arrow points to outer surface of microcapsule membrane. (B) Same biopsy as (A), adjacent section. Antiinsulin immunohistochemistry, showing intensely insulin-positive beta cells, occupying approx. 80% of islet (approx. 240X). Arrow points to outer surface of microcapsule membrane.

Figure 29A-B: (A) Functioning encapsulated neonatal porcine islets, biopsied day 101, from peritoneal cavity of NOD mouse, treated

with CTLA4-Ig for 21 days. Note absence of NOD cell response on outside of membrane and viable islet within capsule (H&E) (X240). Arrows point to outside of capsule wall. (B) Same biopsy as (A). Adjacent section showing majority of cells positive for insulin. Antiinsulin immunocytochemistry (X240).

- 5
- 10 Figure 30: Biopsy of microencapsulated pig islet xenograft from functioning (normoglycemic) transplant into spontaneously diabetic NOD mouse, day #239 after transplantation. Note viable islet cells within microcapsule.
- 15 Hematoxylin & Eosin (H&E), X400. NOD treated with CTLA4Ig for only 21 days after grafting.
- 20 Figure 31: Same biopsy as in Fig. 29; anti-insulin immunohistochemical stain, showing insulin (dark brown) granules in cytoplasm of pig islet "beta" cells within intact microcapsule. Note absence of host NOD cellular reaction on outside of microcapsule membrane (X400).
- 25
- 30 Figure 32: Microencapsulated pig islet xenograft biopsied on day #83 after successful transplant to CTLA4Ig-treated NOD mouse. Note intact "double-walled" microcapsule membrane, viable pig islet cells, and absence of NOD cellular reaction on outside of membrane (H&E, X400).
- 35 Figure 33: Microencapsulated pig islet xenograft

0004965 037060 59864060

5

10

15

20

25

30

35

Figure 40A-B: (A) IL-2 secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) IL-2 secreted by spleen cells (SPC) cultured with porcine

islets, transplanted NODs - rejecting.

Figure 41A-B: (A) IFN-gamma present in peritoneal fluid on sacrifice (sac) day, transplanted NODs.
5 (B) IFN-gamma present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

10 Figure 42A-B: (A) IFN-gamma secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) IFN-gamma secretion by spleen cells (SPC) cultured with porcine islets, transplanted NODs - rejecting.

Figure 43A-B: (A) IL-4 present in peritoneal fluid on sacrifice (sac) day, transplanted NODs.
(B) IL-4 present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

20 Figure 44A-B: (A) IL-4 secreted by spleen cells (SPC)
cultured with porcine islets, transplanted
NODs - not rejecting. (B) IL-4 secreted by
spleen cells (SPC) cultured with porcine
25 islets, transplanted NODs - rejecting.

Figure 45A-B: (A) IL-5 present in peritoneal fluid on sacrifice (sac) day, transplanted NODs.
(B) IL-5 present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

Figure 46A-B: (A) IL-10 present in peritoneal fluid on sacrifice (sac) day, transplanted NODs.
(B) IL-10 present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

5 Figure 47A-B: (A) IL-10 secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) IL-10 secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - rejecting.

10 Figure 48A-B: (A) IL-12 present in peritoneal fluid on sacrifice (sac) day, transplanted NODs. (B) IL-12 present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

15 Figure 49A-B: (A) IL-12 secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) IL-12 secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - rejecting.

20 Figure 50A-B: (A) TNF-alpha (TNF- α) present in peritoneal fluid on sacrifice (sac) day, transplanted NODs. (B) TNF-alpha (TNF- α) present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

25 Figure 51A-B: (A) TNF-alpha (TNF- α) secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) TNF- α secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - rejecting.

30 Figure 52A-B: (A) TGF-beta (TGF- β) present in peritoneal fluid on sacrifice (sac) day, transplanted NODs. (B) TGF-beta (TGF- β) present in peritoneal fluid on sacrifice (sac) day, untransplanted mice.

35

000495E 0279
B6Z2ED 59B54060

5 Figure 53A-B: (A) TGF-beta ($\text{TGF-}\beta$) secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - not rejecting. (B) TGF-beta ($\text{TGF-}\beta$) secreted by spleen cells (SPC) cultured with porcine islets, transplanted NODs - rejecting.

10 Figure 54A-B: (A) NO_2 produced by spleen cells (SPC) cultured with pig islets, transplanted NODs - not rejecting. (B) NO_2 produced by spleen cells (SPC) cultured with pig islets, transplanted NODs - rejecting.

15 Figure 55A-B: (A) NO_2 produced by peritoneal exudate cells (PECs), i.e. from the "belly", after 96 hr in culture with original encapsulated porcine islets, transplanted NODs - not rejecting. (B) NO_2 produced by (PECs) after 96 hr in culture with original encapsulated porcine islets, transplanted NODs - rejecting.

20

25 Figure 56: Proliferation of spleen cells (SPC) from NOD #1335 to pig islets or insulin (Expt 288)

30 Figure 57A-B: (A) Proliferation of spleen cells (SPC) from NOD #1335 to pig islet cells in vitro (Expt 288); (B) Proliferation of spleen cells (SPC) from NOD #1335 (Expt 288).

35 Figure 58: NO_2 Production by cells from NOD #1335 (Expt 288).

0004985 59864060

Detailed Description of the Invention

5 This invention provides a method of inhibiting viable cells
transplanted into a subject from being destroyed by the
subject's immune system which comprises: a) containing the
viable cells, or tissue comprising the viable cells, prior
to transplantation within a device comprising a
semipermeable membrane; and b) treating the subject with a
10 substance which inhibits an immune-system costimulation
event in an amount effective to inhibit the subject's
immune system from responding to said contained cells or
tissue.

15 As used herein, an "immune-system costimulation event" is
an interaction between an APC and a T-cell required in
conjunction with the binding of an MHC-bound antigen on the
surface of the APC to the T cell receptor. As used herein,
APCs are "antigen presenting cells", which are known to
20 those of skill in the art. Immune-system costimulation
events include any specific binding of an APC cell-surface
molecule (other than an MHC-bound antigen) to a specific
ligand on a T cell. Such specific bindings include, but
are not limited to, binding of a B7 molecule (present on
25 the surface of an APC) to a CTLA4 receptor or a CD28
receptor on the surface of a T cell, and binding of a CD40
molecule (present on the surface of an APC) to GP39 (on the
surface of a T cell).

30 Substances which inhibit immune-system costimulation events
are known in the art and include, but are not limited to,
T cell or APC cell-surface-molecule analogs, such as MR1
(which blocks the binding of CD40 expressed on the surface
of an APC to GP39 expressed on the surface of a T cell), or
35 CTLA4 (which blocks the binding of a B7 molecule to a CD28

receptor or a CTLA4 receptor).

8622E0"59864060

In one embodiment of the method for inhibiting destruction of viable transplanted cells described herein, the substance which inhibits an immune-system costimulation event is CTLA4. The term CTLA4, for purposes of this invention, is meant to indicate any proteinaceous construct which comprises an amino acid sequence which is the same as or sufficiently the same as the amino acid sequence of the CTLA4 receptor such that the proteinaceous construct is capable of binding to a B7 molecule, thereby blocking the B7 molecule from binding to a CTLA4 receptor on a T cell. Proteinaceous constructs are well known in the art and indicate any molecule which comprises amino acid moieties linked to one another by peptide bonds; including peptides, polypeptides, and molecules comprising peptide and/or peptide subunits. Thus, the term CTLA4 includes, but is not limited to, molecules expressed by the gene encoding the B7-binding site of the CTLA4 receptor in genetically engineered cells, molecules expressed by mutants of the gene encoding the B7-binding site of the CTLA4 receptor which molecules are capable of binding to a B7 molecule, and synthetic amino acid chains having an amino acid sequence which is the same as or sufficiently the same as the amino acid sequence of the CTLA4 receptor such that they are able to bind to B7. CTLA4 also includes soluble CTLA4 comprising the extracellular binding domain of the CTLA4 receptor, such as CTLA4Ig. Accordingly, the term CTLA4 for purposes of this invention also includes CTLA4Ig, i.e. a recombinant soluble fusion protein which combines the extracellular binding domain of the CTLA4 receptor with the constant region of IgG₁.

In an embodiment of this invention, the substance which inhibits an immune-system costimulation event also alters

15 In another embodiment, the substance which inhibits an
immune-system costimulation event binds complement.
Substances which bind complement favor prolonged survival
of contained cells or tissue grafted into the subject. An
20 example of a substance which binds complement is CTLA4Ig.

In another embodiment of this invention, the substance which inhibits an immune-system costimulation event does not alter the cytokine profile of the subject so as to protect the contained cells or tissue from the subject's immune system. The term "cytokine profile" means the type and quantity of each type of cytokine produced in a subject at a given time. Cytokines are proteins which have an immune effect and which are released by white blood cells. Examples of cytokines include, but are not limited to interferon (such as gamma-interferon), tumor necrosis factor, interleukin (IL) 1, IL-2, IL-4, IL-6, and IL-10. For example, the substance may be a substance which increases the production of gamma-interferon and IL-2 in the subject. An example of a substance which does not

alter the cytokine profile of a subject so as to protect contained cells or tissue grafted into the subject is CTLA4Ig. In an embodiment, the substance and the containing of the viable cells within the device comprising
5 the semipermeable membrane prevents host immune cell proliferation in the subject. In an embodiment of the above-described method the device comprising the semipermeable membrane is a hollow, fiber, a disc, or a sphere. In a further embodiment of the above-described
10 method the device comprising the semipermeable membrane is a microcapsule.

This invention also provides a method of inhibiting viable cells transplanted into a subject from being destroyed by
15 the subject's immune system which comprises: a) containing the viable cells, or tissue comprising the viable cells, prior to transplantation within a device comprising a semipermeable membrane; and b) treating the subject with CTLA4 in an amount effective to inhibit the subject's
20 immune system from responding to said contained cells or tissue.

Devices comprising a semipermeable membrane useful for transplantation of viable cells or tissue are well-known to
25 those of ordinary skill in the art, and any such device may be used in the subject invention. Devices useful for the subject invention may be comprised of various materials and may be formed into various shapes, such materials and shapes being well known in the art. Any particular device
30 for an application of this invention is selectable based on factors including, but not limited to, the biocompatibility of the material with the subject, the site of transplantation, whether the transplantation is intravascular or extravascular, the method of
35 transplantation, availability, and economy. Examples of

85/23E9 59864060

5

10

20

35

5

10

15

30

encoding the hormone and are capable of expressing the gene.

5 In the aforementioned method of this invention, the viable cells in one embodiment comprise cells which secrete a biologically active substance. The term "biologically active substance" as used herein means any substance which is capable of eliciting a physiological response in a subject. The biologically active substance may illicit a response in the subject into which the cells producing it are transplanted. Cells which secrete biologically active substances are well known in the art, and any such cells may be used in the subject invention.

10

15 In one embodiment, the cells which secrete a biologically active substance are endocrine cells. Endocrine cells are well known to those of ordinary skill in the art and include, but are not limited to, insulin-producing cells, hepatocytes, parathyroid cells, and pituitary cells. In another embodiment, the cells which secrete a biologically active substance are neuroectodermal cells. Neuroectodermal cells are also well known in the art, and include, but are not limited to, adrenal cells and lymphocytes.

20

25 In another embodiment, the cells are genetically engineered to secrete a biologically active substance. For example, the cells may be genetically engineered to secrete a biologically active substance useful for treating the subject into which they are transplanted. Thus, the subject method provides a novel, useful, and advantageous drug delivery system for treatment of subjects afflicted with conditions including, but not limited to, cancer and HIV infection. If the subject is afflicted with cancer,

30

35 the transplanted viable cells may, for example, be

004985 032798
862220 59864060

genetically engineered to secrete Interleukin-2, a cytokine, or a lymphokine. If the subject is infected with HIV, the transplanted viable cells may, for example, be genetically engineered to secrete a substance which stimulates lymphocyte production in the subject, such as a T cell growth factor or the HIV T cell receptor.

In the method of the subject invention, the permeability of the semipermeable membrane of the device is determined based on factors well known in the art, for example, the size of the cells or tissue being contained, the size of any substances needed to permeate the membrane in order to sustain the cells or tissue, and the size of any biologically active substances secreted by the cells which are desired to permeate from the device. In one embodiment, the semipermeable membrane is impermeable to lymphocytes. In another embodiment, the semipermeable membrane is impermeable to lymphocytes and immunoglobulins. Using a semipermeable membrane which is impermeable to immunoglobulins and/or lymphocytes prevents contact between the immunoglobulins and/or lymphocytes of the subject and the contained viable cells, and thereby prevents destruction of the contained cells which would result from such contact.

Any suitable method of treatment may be used in the subject invention to treat the subject with the substance which inhibits an immune-system costimulation event, and such methods are well-known in the art. For example, the substance may be administered by injection to the subject in the form of a pharmaceutically acceptable composition. If the substance is CTLA4, CTLA4Ig may be directly administered to the subject, or in another embodiment, cells genetically engineered to secrete CTLA4, that is cells which have been genetically engineered to contain a

gene encoding a molecule capable of binding to a B7 molecule and to express that molecule, may be transplanted into the subject.

5 In another embodiment of the invention, treatment of the subject with the substance comprises transplanting into the subject cells genetically engineered to secrete the substance. If cells genetically engineered to secrete the substance are transplanted into the subject, such cells may
10 themselves be contained within a device comprising a semipermeable membrane prior to transplantation. In different embodiments, the semipermeable membrane of the device containing the cells secreting the substance is impermeable to immunoglobulins and/or lymphocytes, thereby
15 preventing destruction of these cells which would otherwise result from such contact.

In the aforementioned embodiments, treatment with the substance may occur before, after, of contemporaneously
20 with transplantation of the viable cells or tissue.

In another embodiment of the subject invention, treating the subject with the substance comprises containing cells genetically engineered to secrete the substance within the
25 device containing the viable cells or tissue prior to transplantation.

In a further embodiment of the invention, treating the subject with the substance comprises genetically
30 engineering the viable cells transplanted into the subject to secrete the substance prior to transplantation.

The amount of the substance effective to inhibit the subject's immune system from responding to said contained
35 cells or tissue is determined by factors well-known to

854220 59864060

those of skill in the art, including, but not limited to, the amount of viable cells or tissue transplanted into the subject, and the size and weight of the subject.

- 5 Inhibiting the subject's immune system from responding to the contained viable cells or tissue by the method of the subject invention involves an inhibition of immunoglobulin production in the subject and an inhibition of macrophage activation in the subject. Such immunoglobulins and
10 activated macrophages would otherwise be capable of reacting with, and destroying, the contained viable cells or tissue.

- This invention also provides a method of treating diabetes
15 in a subject which comprises: a) containing viable insulin-producing cells, or tissue comprising viable insulin-producing cells, within a device comprising a semipermeable membrane so as to obtain contained viable insulin-producing cells; b) transplanting contained viable insulin-producing
20 cells obtained in step (a) into the subject in an amount effective to treat diabetes in the subject; and c) treating the subject with a substance which inhibits an immune-system costimulation event in an amount effective to inhibit the subject's immune system from responding to an
25 amount of contained viable insulin-producing cells according to step (b).

- Substances which inhibit an immune-system costimulation event are known in the art, and any such substance may be
30 used in the method for treating diabetes described herein. Substances which inhibit an immune-system costimulation event which may be used in the subject method for treating diabetes are described above. In one embodiment, the substance is CTLA4.

The viable insulin-producing cells, or tissue comprising viable insulin-producing cells, may be obtained from any known source for insulin-producing cells or tissue comprising insulin-producing cells.

5

In one embodiment of the subject invention, viable insulin-producing cells are derived from pancreatic islet tissue. In another embodiment, the viable insulin-producing cells comprise cells which have been genetically engineered prior to transplantation to secrete insulin. The viable cells or tissue may be derived from a xenogeneic donor, an allogeneic donor, or they may be derived from the subject prior to transplantation. If the cells are derived from the subject, in one embodiment, they are genetically engineered to produce insulin after they have been removed from the subject, prior to being transplanted back into the subject.

The viable insulin-producing cells or tissue comprising viable insulin-producing cells, such as a pancreatic islet tissue, may be obtained from any donor. In one embodiment, the donor is a mammal. Such a mammalian donor may, for example, be a calve, a pig, a rabbit, a rat, a mouse, or a human. The viable insulin-producing cells or tissue comprising viable insulin-producing cells, such as pancreatic islet tissue, may be obtained from a mammalian neonate, such as a neonatal pig. In one embodiment, the viable insulin-producing cells or tissue comprising viable insulin-producing cells used in the subject invention comprises neonatal porcine (pig) pancreatic cells.

The subject of the invented method described herein may be any subject into which transplantation of viable cells is desired. In one embodiment, the subject is a human. If the subject is a human, the viable cells, or tissue

containing them, are in one embodiment derived from a mammal, for example a human.

5 Devices comprising a semipermeable membrane are well-known to those of ordinary skill as described above, and any such device may be used in the subject method of treating diabetes. In different embodiments of the method, the device is a hollow fiber, a disk, and a sphere. In another embodiment of the method, the device is a microcapsule as
10 described above.

The method of treating diabetes described herein may be applied to any subject for whom diabetes treatment is desired. In one embodiment of the invented method for
15 treating diabetes in a subject, the subject is afflicted with insulin-dependent diabetes mellitus (IDDM). In another embodiment of the method, the subject is a mammal, for example a human.

20 The amount of contained viable insulin-producing cells transplanted into the subject effective to treat diabetes in the subject depends on factors known to those of ordinary skill, including, but not limited to, factors such as the weight of the subject, and the severity of the
25 diabetes.

The permeability of the semipermeable membrane of the device in the subject method of treating diabetes is determined by factors known to those of ordinary skill,
30 including those factors for determining permeability described above. In different embodiments of the method, the semipermeable membrane is impermeable to immunoglobulins and/or lymphocytes.

35 Treatment of the subject with the substance which inhibits

552220 59867060

In the aforementioned methods of treating the subject with a substance, such as CTLA4, capable of inhibiting an immune-system costimulation event, treatment may occur before, after, or contemporaneously with transplantation of the contained viable insulin-producing cells into the subject.

In another embodiment of the subject method of treating diabetes, treating the subject with the substance comprises 35 genetically engineering the viable insulin-producing cells

Inhibiting the subject's immune system from responding to contained viable insulin-producing cells or tissue by the subject method of treating diabetes involves an inhibition of immunoglobulin production and of macrophage activation in the subject which would otherwise react with and lead to the destruction of the viable insulin-producing cells or tissue.

This invention will be better understood from the "Experimental Details" section which follows. However, one skilled in the art will readily appreciate that the specific methods and results discussed therein are not
15 intended to limit, and rather merely illustrate, the invention as described more fully in the claims which follow thereafter.

Experimental Details

Improvements in Microcapsule Design

An improved formulation of poly-L-lysine-alginate microencapsulation which allows nearly indefinite survival of rat islets in spontaneously diabetic NOD mice is the "double-wall" microcapsule (Figures 1 and 2). This double-wall microcapsule is more durable than conventional microcapsules, with fewer capsule wall defects, has a measured membrane permeability of approximately 100,000 Kd, and excludes IgG (unlike conventional design capsules, which allowed passage of IgG and 148,000 Kd fluoresceinated dextran) (9,19,20,118). These data support the relevance of encapsulated islet xenografts for eventual application in humans with IDDM.

Poly-L-Lysine (PLL) Concentration Alters Permeability of PLL-Alginate Microcapsules

It was postulated that microencapsulated islet xenograft survival would be influenced by microcapsule permeability. We found that microcapsule permeability may be altered by increasing or decreasing the concentration of PLL (poly-L-lysine) in the microcapsule formula. Red blood cells were encapsulated in alginate via an air jet system and then incubated with various polyamino acids including PLL. The RBCs were then lysed and hemoglobin (MW 64,500) efflux was measured spectrophotometrically at 480nm as a function of time alongside a concurrent control. Permeability coefficient was calculated according to the following formula: $(2.303 * C_f * V_t * S) / (C_i * A_t)$, where C_i and C_f are the initial and final hemoglobin concentrations, V_t and A_t are the total volumes and areas of capsules respectively, and $S = \text{slope of } \ln (C_t - C_f) / (C_i - C_t)$ (119). PLL substitutions (poly-L-ornithine, alanine, aspartate and histidine) did not result in viable capsules. PLL molecular weight alterations did not effect permeability.

PLL concentration was the most critical factor in altering capsule diffusion. These observations are supported by the recent findings of other investigators (119). There was a thirteen fold decrease in hemoglobin efflux occurring
5 in capsules that had a fourfold increase in PLL (see Table 1). In experiments, encapsulated rabbit islet survival in NODs is prolonged using microcapsules with permeability <70,000 Kd vs. 100,000 Kd (see Figure 3).

05/22/00 5:58:40 PM

Table 1 Increasing PLL Concentration Reduces Microcapsule Permeability to Hemoglobin

PLL Concentration (% w/v)	0.050	0.125	0.137	0.144	0.150	0.200
Permeability constant (E-06cm/sec)	50	56	52	30	6.7	3.8

Microcapsules Prevent or Delay Host Sensitization

To clarify the mechanism of long-term microcapsule protection of xenogeneic rat islets, experiments were performed in which paired diabetic NODs were pre-treated with saline or Lewis rat islets (200 intra-peritoneally) or 10^6 Lewis rat splenocytes intra-peritoneally. Encapsulated Lewis islets were xenografted into presensitized and control NODs 14 days later. As shown in Figures 4 and 5, both islet- and splenocyte pretreatment resulted in rapid graft rejection while non-presensitized NODs accepted encapsulated islet xenografts long-term. These data suggest that a major function of microcapsules is to prevent host sensitization, rather than to protect grafts from the effector arm of the response. Thus, maneuvers which reduce islet Immunogenicity may be synergistic with islet encapsulation.

Comparisons of Encapsulated Islet Iso-, Allo- and Xenograft Survival in NODs

We have found that microencapsulation allowed islet xenograft survival in NODs of 79 ± 15 days ($N=8$) ($X \pm SE$) for Lewis rat islets, vs. 20 ± 2 days ($N=7$) for rabbit islets and 14 ± 4 ($N=3$) for dog islets (Table 2), with similar peri-microcapsule NOD cell accumulations at rejection. NODs also rejected encapsulated, allogenic Balb/c islets in 73 ± 31 days ($N=4$) and encapsulated isologous NOD islets in 44 ± 7 days ($N=4$) (Table 2). However, biopsies of these allo- and isologous grafts, at rejection, have shown few host macrophages adherent to microcapsules, while free peritoneal cells (thus far not characterized) were present. Thus encapsulated islet xenograft rejection is distinct from iso- and allo-graft rejection in this model.

Table 2.

Islet Iso-, Allo- and Xenografts in NOD Mice

*=P<.002 vs. Group 7; @= Mean \pm SEM; #=P<.05 vs. Group 7;

^{**}=P<.003 vs. Group 7

CAP/I.P. = microencapsulated islet graft to peritoneal cavity;
Splenic = Nonencapsulated islets grafted beneath splenic
capsule.

We have also found that microencapsulation prolongs the functional survival of islet xenografts in NODs, when compared to survival of unencapsulated islets injected

into the spleen. The same is true for islet allografts and for islet isografts into NODs (Table 3).

Table 3.

Beneficial Effect of "Double-Wall" Microencapsulation of Survival of Islet Iso-, Allo-and Xenografts in NOD Mice

Donor-Recip	Technique	(N)	Surv(days)@
NOD-NOD	CAP/I.P.	4	44±7*
NOD-NOD	Splenic	3	6,7,7
Balb-NOD	CAP/I.P.	4	73±31*
Balb-NOD	Splenic	2	5,5
Lewis Rat-NOD	CAP/I.P.	8	79±15*
Lewis Rat-NOD	Splenic	9	19±3
Dog-NOD	CAP/I.P.	3	14±4*
Dog-NOD	Splenic	2	0,0
Rabbit-NOD	CAP/I.P.	7	20±2*
Rabbit-NOD	Splenic	2	5,6
Neonatal Pig-NOD	CAP/I.P.	8	27±13*
Neonatal Pig-NOD	Splenic	3	6±1

p<.01 vs. splenic: @=Mean ± SEM.CAP/I.P. = microencapsulated islet graft to peritoneal cavity;
Splenic = Nonencapsulated islets grafted beneath splenic capsule.

Functioning and rejected encapsulated xenografts were biopsied from the peritoneal graft sites of spontaneously diabetic NOD mice, on days #4-#50 post-transplantation.

- Controls included normal mouse peritoneal fluid and peritoneal fluid from NOD mice bearing empty capsules or capsules with functioning (recipient normoglycemic) rat islets (20,74). However, cell number increased dramatically at rejection on days #14 and #50. Pipetting

00220"59864060

of biopsied capsules freed adherent cells. Flow cytometric analyses revealed that 20-50% of non-adherent peritoneal cells were B220⁺ (B cells), and that the majority of free peritoneal cells and cells adherent to microcapsules were Mac1⁺ (20,74). The percentages of CD4⁺ and CD8⁺ peritoneal cells were low (4-9%). By FACS analysis, the phenotype of peritoneal Mac1 cells shifted from predominantly Gran1⁻ to Gran 1⁺ during rejection of xenogeneic islets in microcapsules (vs. empty capsules) (20,74,120). These findings were confirmed by immunocytochemistry (20,74). In addition, immunocytochemistry documented IgG and IgM around microcapsules, and IL-1 and TNF alpha both around and within microcapsules (20,74).

15

Analysis of Cytokine Messenger RNA (mRNA) in Encapsulated Islet Xenografts Biopsies from NODs

To elucidate the pathogenesis of NOD destruction of encapsulated islets, mRNA was extracted from recipient NOD peritoneal cells and expression of mRNA for IL-2, IL-4, and IL-10 was studied by RT-PCR, as previously described (121). Integrity of RNA samples was assessed by inspection of northern transfer and hybridization with the probe for the 3' untranslated region of beta actin (121). IL-4 was detected in the majority of xenografts undergoing rejection. IL-10 expression was variable (Table 4). IL-2 was detected during autoimmune destruction of NOD isografts, (and in one allograft) but only rarely in rejecting xenografts (Table 4). These data suggest that the primary T cell response in rejecting encapsulated islet xenografts is "Th2-like". This interpretation is consistent with the observation that large numbers of activated macrophages and immunoglobulins are associated with rejecting encapsulated islet xenografts in NODs. Thus, it is

004965-0329
86220-59864060

possible that rejection of encapsulated islet xenografts is initiated by soluble, or shed, xenoantigens that are processed via the Class II pathway by host APC. These APC then activate Th2 cells via B7/CD28 dependent mechanisms. We postulate that formation of antigen-antibody complexes in the peritoneal cavity activates macrophages to release cytokines that are directly toxic to encapsulated islets.

85220 59864060

Table 4.

5

CYTOKINE mRNA IN BIOPSIES OF ENCAPSULATED XENO- ISLETS IN
NOD MICE

						mRNAs		
	Islet Donor	NOD#	Sample	Day Rejected	Day Biopsied	IL2	IL4	IL10
10	NOD	194	FC*	39	40	+	+	-
		291	FC	14	21	+	-	-
	Balb/c	487	Cap	12	14	+	-	-
15	Rat	154	Cap	18	20	-	+	-
		154	FC	18	20	-	+	-
		58	Cap	34	38	-	+	+
		165	Cap	21	28	-	+	-
		54	Cap	136	143	+	+	+
20		54	FC	136	143	+	-	-
		107	FC	41	45	-	-	-
		453	Cap	132	134	+	-	-
	Canine	141	Cap	17	24	-	+	+
		268	Cap	13	14	-	-	-
25		268	FC	13	14	-	-	-
		69	FC	18	24	-	+	+
	Rabbit	91	Cap	35	49	-	-	-
		91	FC	35	49	-	+	-
		151	Cap	28	32	+	+	-
		46	FC	12	15	-	+	-
		55	FC	18	21	-	+	-
30		152	FC	Funct.	15	-	+	-
		157	FC	Funct.	15	+	-	-
35	Human	136	Cap	6	8	-	+	+

Cap = Cells adherent to capsules

*F C = Free peritoneal cells

† = RT-PCR (-) is undetectable and (+) is detectable

40

45

862220 59854060

The NOD-MHC is Necessary for Rejection of Encapsulated Islet Xenografts

Both NOD and (SZN-diabetic) B10.H-2^{g7} (expresses the NOD-MHC-linked disease allele) rejected encapsulated rat islets, while NOD.H-2^b mice, which express all of the non-MHC-linked diabetes susceptibility genes, accepted encapsulated rat islets for >100 days (similar to B10 controls) (75). This suggests that the NOD-MHC may contribute to destructive responses against encapsulated islets which are distinct from diabetes susceptibility, since neither B10.H-2^{g7} nor NOD.4-2^b mice develop diabetes spontaneously (20,75). The possibility that SZN treatment of B10.H-2^{g7} mice may have initiated an autoimmune response was considered; however, 2/2 non-diabetic (no SZN treatment) B10.H-2^{g7} mice rejected encapsulated rat islets (by biopsy histology, day #60) (75).

CD8⁺ Depletion Does Not Protect Encapsulated Islet Xenografts in NODs

It was found that treatment of NOD recipients of encapsulated rabbit islets with either monoclonal antibody 53.6.7, (100 μ g i.p. day -5 and then twice weekly) (anti-CD8) or cyclosporine (CyA), 30. Mg/kg, s.c., daily had no effect on graft survival (Table 2). CD8⁺ cell depletion was confirmed by flow cytometry of NOD spleen and peritoneal cells. Biopsies of failed grafts revealed intense host cellular responses and non-viable islets within intact microcapsules. These data are consistent with prior observations, that CD4⁺ (but not CD8) T-cells play a dominate role in non-encapsulated islet xenograft rejection (83). They also are consistent with a predominantly Th2 NOD rejection mechanism of encapsulated islet xenografts.

Co-stimulatory Blockade Prolongs Encapsulated Islets Xenografts in Diabetic NODs

It was shown previously that inhibition of CD4⁺ helper T-cells by administration of monoclonal antibody GK 1.5 to diabetic NOD recipients resulted in significantly increased survival (>100 days) of both encapsulated rat and dog islets (7,84) (Figure 6). The experiments herein show that treatment of NOD mice with CTLA4Ig (200µg i.p. day #0, and QOD until day #90) significantly prolonged encapsulated rabbit islet survival, from 20 ± 2 days to 98 ± 25 days (p<.05) (see Table 2 and Figures 7 and 8).

10

This suggests that an "indirect" pathway of antigen presentation is dominant in NOD responses to encapsulated islet xenografts. Unlike findings with human islet transplanted to SZN-diabetic mice (12), CTLA4Ig alone did not increase nonencapsulated rabbit or rat islet survival in NODs (intrasplenic or renal subcapsule) (Table 2), suggesting that encapsulation and CTLA4Ig both were required to prolong graft survival.

20 Furthermore, the experiments herein show that encapsulated female islets from INSCTLA4 mice, which express CTLA4 on the beta cell insulin promoter, function long-term in NODs (see Figure 9). Unencapsulated INSCTLA4 islets were rejected by NODs in 6-7 days. These data suggest that indefinite survival of discordant islet xenografts may be achieved by combinations of donor islet encapsulation and limited host immunomodulation. These data also support the working hypothesis that donor antigen(s) are shed from microcapsules and processed by APCs which activate CD4⁺ T cells via B7/CD28-dependent mechanisms. In this model, CTLA4-transgenic mice secrete CTLA4, along with insulin, and CTLA4 inhibits antigen presentation. Interestingly, female mice secrete more CTLA4 than do male mice in this transgenic model (pers. Comm.).

35

NOD-Scid Mice Accept Rat and Rabbit Islet Xenografts Long-Term

These experiments demonstrate that NOD-scid mice are susceptible to MLD-SZN diabetes (30mg/kg daily x5); and reversal of NOD-scid diabetes with xenografts of nonencapsulated and encapsulated rat and rabbit islets for greater than 50 days is documented (see Figures 10,11, and 12 and Table 2). Thus, the NOD-scid mice will serve as a good recipient model for the transfer of antibodies and/or T cells for studies of the mechanisms by which encapsulated islets are rejected. We noted recurrent hyperglycemia in 3/4 NOD-scids receiving microencapsulated rabbit islets, on days #51, #68, and #70. Biopsies revealed disrupted capsules and minimal cellular failure for technical reasons, since empty microcapsule controls done concurrently, showed broken microcapsules (in 1/3) and intact microcapsules (in 2/3) at day #50.

Costimulation Blockade with CTLA4Ig

20 Method:

Adult New Zealand rabbit islets were isolated by duct-injection, collagenase digestion. Rabbit islets (approx. 2000) were encapsulated in double-wall, poly-L-lysine-alginate microcapsules and xenografted intraperitoneally in NODs, as previously reported (7,20). Controls received approximately 2000 unencapsulated rabbit islets xenografted beneath the splenic or renal capsule, as previously described (7,20).

30 Murine CTLA4Ig, provided by Bristol-Myers-Squibb, Seattle, WA, was administered at 200ug intraperitoneally (i.p.), day-1 and then Q.O.D. for 14 or 92 days, or until graft rejection.

35 Controls included NODs receiving identically encapsulated

004986 03298

5

10

15

20

25

30

35

Table 5:

EFFECTS OF CTLA4Ig, CyA AND ANTI-CD8
MONOCLONAL ANTIBODY ON ENCAPSULATED RABBIT
ISLET XENOGRAFT SURVIVAL IN DIABETIC NOD MICE

Group	Donor-Recip	Technique	Rx.	(N)	X±SE	Graft Survival
						Days
#1	Rabbit-NOD	CAP/I.P.	None	7	20±2	12, 16, 18, 18, 20, 28, 28
#2	Rabbit-NOD	CAP/I.P.	CyA	4	22±3	13, 24, 26, 26
#3	Rabbit-NOD	CAP/I.P.	53.6.7.7 7.7	4	5±9	14, 15, 18, 52
#4	Rabbit-NOD	CAP/I.P.	CTLA4Ig (x92 days)	8	108±24 ⁺	37 ^d , 43, 47, 58 148, 151 ^s , 173, 205 ^d
#5	Rabbit-NOD	CAP/I.P.	CTLA4Ig (x14 days)	4	70±8 ^{**@}	48, 66, 81, 83
#6	Rabbit-NOD	Renal/ Splenic	CTLA4Ig	3	6±1 [*]	5 ^(s) , 6 ^(r) , 6 ^(a)
#7	Rabbit-NOD	Renal/ Splenic	None	2	-	5 ^(a) , 6 ^(r)

s = sacrificed, functioning graft.

d = died, functioning graft.

*p<.005 vs. Group 1, ("t"-test).

**p<.0001 vs. Group 1, ("t"-test).

CTLA4Ig, 200µg day -1, then Q.O.D., i.p.

CyA - 30mg/kg day -1, then Q.D., s.c.

63.6.7.7 -100µg, day -5,+2, then weekly, i.p.

(r) = renal subcapsule, not encapsulated

(s) = splenic subcapsule

@ = P = .31 vs. Group #4, ANOVA

Large-Scale Neonatal Porcine Islet Isolation

We believe the neonatal pig is the most promising xenogeneic source of donor islets. A reproducible method for isolation of large numbers of functionally viable islets from neonatal porcine donors has been developed (146,147). With this technique, 30,000-100,000 islets may
5 be obtained from each donor pig (Figure 14). Neonatal pig islet cells continue to secrete insulin *in vitro* after microencapsulation. (Figure 15). These neonatal pig islets are actually dispersed neonatal porcine pancreatic cells which reaggregate to form "islet"-like spheroids with
10 approximately 5-10% beta cells (Figure 16), which is significantly higher than the 1-2% beta cell concentration in the adult porcine pancreas. Furthermore, biopsies of these "islets" 100 days following xenotransplantation reveal increased numbers of intensely insulin-positive
15 islet cells (Figure 17). These neonatal pig islets have an added advantage over adult islets, in that they appear to differentiate and proliferate within microcapsules after transplantation.

20 **Both Encapsulated and Non-encapsulated Neonatal Porcine Islets Reverse SZN-Diabetes in NOD-Scid Mice.**

Recently, the Scid mutation has been back-crossed onto the NOD background, resulting in immuno-deficient NOD-Scid mice (66,67,68,69). These mice are homozygous for the Scid
25 mutation, which results in an inability to rearrange T-cell receptor and immunoglobulin genes (48,79). Consequently, these mice lack T and B-lymphocytes. NOD-Scid mice do not develop diabetes spontaneously; but they may be rendered diabetic with multiple low-dose streptozotocin (MLD-SZN),
30 (67,68,69) NOD-Scids express NOD MHC genes and other genes that are required for development of diabetes, upon transfer of lymphocytes from diabetic NODs.

To document functional viability of neonatal porcine

0004985-032798

Table 6. Survival of Microencapsulated (MC) Adult Rabbit and Neonatal Procine Islets in NOD Mice:
Effects of NOD Treatment with CTLA4Ig

Graft Survival					
Donor	Technique	Rx.	(N)	x±S.E.	Days
Rabbit	MC/I.P.	None	7	20±2	12, 16, 18, 18, 20, 28, 28
Rabbit	MC/I.P.	CTLA4-Ig ^a	8	108±24*	37, 43, 47, 58, 148, 151, 173, 205
Rabbit	MC/I.P.	CTLA4-Ig ^b	4	70±8*	48, 66, 81, 83
Rabbit	Splenic	CTLA4-Ig ^a	3	6±1*	5, 6, 6
Rabbit	Splenic	None	2	--	5, 6
Neonatal Pig	MC/IP	CTLA4Ig ^c	8	27±13*	9, 10, 12, 12, 14, 14, 23, 118 ^(s)
Neonatal Pig	MC/IP		5	111±17*	74 ^(s) , 80, 101 ^(s) , 137 ^(s) , 161 ^(s)
Neonatal Pig	Splenic	None	3	5±1	4, 5, 5
Neonatal Pig	Splenic		3	6	5, 6, 7

IP=intraperitoneal

CTLA4Ig, 200 mcg I.P., QOD

* = p >.001 vs. MC alone

- a) x92 days
- b) x14 days
- c) x21 days

(s) = sacrifice for biopsy

Biopsies of long-term functioning encapsulated neonatal porcine islet xenografts showed viable porcine islets within intact microcapsules and absence of host NOD pericapsular reactivity was observed in biopsies of long-term normoglycemic NODs (Figure 22 and 23).

To analyze the potential mechanisms of action of CTLA4Ig in this model, we substituted a recently devised mutant of CTLA4Ig, which does not fix complement (CTLA4Ig*) (145). As shown in Figure 24, our studies have revealed that CTLA4Ig* does not prolong graft survival above that of capsules alone. The data are distinct from findings with murine allografts, which are prolonged significantly by either conventional CTLA4Ig or mutant CTLA4Ig*. These results suggest that mechanisms of prolongation of graft survival by CTLA4Ig* may be different for allogeneic and xenogeneic islet grafts. The results suggest that the cytokine profile in a subject can be altered in favor of graft protection. In the system studied in this experiment, conventional CTLA4Ig altered the cytokine production so as to protect the graft by increasing gamma-interferon production in the host. Conversely, in the studied system, an increase in IL-10 production induced by CTLA4Ig* treatment favored graft rejection.

We also measured proliferative responses by spleen cells from a matched pair of diabetic NOD mice that were transplanted with the same batch of encapsulated, neonatal pig islets but were treated with either CTLA4Ig or the non-complement fixing CTLA4Ig* (Figure 25). In this experiment, normal or diabetic NOD mice did not proliferate when stimulated by neonatal pig islets (panel A and B). The reason for the inconsistent response of nontransplanted NOD mice is not yet known but is under investigation. Empty capsules did not induce proliferation in any of the

spleen cells but islets and encapsulated islets recognized by T-cells are small enough to exit from microcapsules. However, more experiments may verify this interpretation. As usual, background responses of spleen cells from mice rejecting grafts (panel D) were higher than those from mice that were not rejecting grafts (panel C).

These results suggest that spleen cells from both mice engrafted with encapsulated islets were primed *in vivo*, and are somewhat surprising given the fact that the mouse that received CTLA4Ig showed no signs of rejection. These results did not address the possibility that there might be different fluids from cultures stimulated with neonatal pig islets for lymphokines by ELISA (Figure 26). These results indicate that lymphokines were produced only by mice that were engrafted with neonatal, pig islets. More importantly, spleen cells from the mouse that had accepted its graft long term (treated with CTLA4Ig) produced a preponderance of $\text{INF}\gamma$ and low levels of IL-10. These results suggest that CTLA4Ig induced long term tolerance to neonatal pig islets that is associated with T cells that produce $\text{INF}\gamma$. Rejection of xenogeneic islet graft occurred when lymphokines shifted to IL-10. Thus, graft rejection is associated with a Th2-like response, whereas graft survival is associated with Th1-like responses. These findings are consistent with our working model (Figure 27). These results differ somewhat from the picture obtained by analyzing mRNA level at the site of rejection where IL-4 predominated in mice that rejected the encapsulated, xenogeneic islets.

Discussion:

On the basis of our data, we develop a model to describe the mechanisms that we think are involved in rejection of microencapsulated xenogeneic islets by autoimmune, NOD mice

(Figure 27). Secreted insulin clearly crosses the membrane of double walled microcapsules and regulates glucose levels in engrafted mice. Potentially, other donor proteins or protein fragments of less than 100,000 mw (AgX) that are shed or secreted by islets diffuse out of the microcapsule and are endocytosed by dendritic cells. Dendritic cells process proteins via the MHC class II pathway and present peptide X complexed with class II and co-stimulatory molecules to CD4⁺T cells. In the presence of the appropriate cytokines, CD4⁺T cells are activated and develop into Th2 cells that express CD40L. B cells with surface IgM that binds AgX endocytose and process it into peptides that bind MHC class II which are expressed on the surface of B cells. Th2 specific peptide X complexed with class II binds B cells and the interaction of CD40 with CD40L causes the activation of B cells. Activated B cells mature into plasma cells under the direction of Th2 lymphokines. Plasma cells secrete specific antibody that forms complexes with AgX.

Antibodies are not able to directly damage the encapsulated islets because they are too large to enter the capsules. However, antibodies could be involved in the recruitment and activation of macrophages which are the predominant population in the peritoneal cavity of NODs rejecting encapsulated islet xenografts. Specific antibodies in the peritoneal cavity could form complexes with antigens shed or secreted from the capsules. Such antigen-antibody complexes efficiently bind to FcR expressed on the surface of peritoneal macrophages. Binding of complexes to FcR activates macrophages to secrete a variety of mediators including IL-1, TNF α and nitric oxide (NO) (122,123), all of which have toxic effects on islets and all of which are small enough to cross a double walled microcapsule. The effector arm could be further augmented by the activation

of complement (c) by antigen complexes. C3b bound to the complexes enhances the activation of macrophages by increasing the binding of the complexes via the C3b receptor (124) and small peptides such as C3b released during complement activation induce local inflammatory responses thereby attracting more macrophages into the peritoneal cavity (125).

We demonstrated synergy of donor islet microencapsulation and NOD CTLA4Ig treatment in prolonging islet xenograft survival. Our data represent the longest biopsy-proven survival of discordant islet xenografts in NODs reported to date. Neither CTLA4Ig nor encapsulation alone were effective. Furthermore, splenocytes from a long-term successful graft recipient did not transfer donor-specific unresponsiveness. Failure of anti-CD8 and CyA therapies is consistent with our hypothesis of a primarily Th2 type response in this model.

There is considerable evidence that xeno-recognition (unlike allorecognition) occurs primarily via the so-called "indirect" antigen presentation pathway, by which host APC present peptides scavenged from extracellular (donor) proteins to host helper T-cells (27,137,29,138). Our recent report, that the host MHC is critical to NOD rejection of encapsulated islet xenografts (75), and our prior observations, that helper T-cells are essential for this response (7), both are consistent with an "indirect" pathway. Our prior findings of more rapid destruction of encapsulated "discordant" (widely unrelated) islets (canine, rabbit, bovine, porcine) than "concordant" (closely related)(rat) islets (20), also support this hypothesis, since the "direct" pathway would favor an accelerated reaction to "concordant" donor tissue. Furthermore, our current data suggest that "indirect"

antigen presentation may be blocked by CTLA4Ig in this model of encapsulated islet xenotransplantation. In conclusion, we have found that neither microencapsulation nor CTLA4Ig alone prevent NOD destruction of rabbit islets.

5 However, we have observed synergy between CTLA4Ig treatment
of NOD recipients plus encapsulation with significantly
prolonged discordant islet xenograft survival.

References

1. The Diabetes Control and Complications Trials Research Group. The effect of intensive treatment of diabetes on the development and progressions of long-term complications in insulin-dependent diabetes mellitus. NEJM 1993;329:977-986.
2. Lacy P. Status of islet cell transplantation. Diabetes Review 1993;1:76-92.
3. Parker C. Naji A. Perspectives and islet transplantation for diabetes-cures or curiosities? NEJM 1992;327:1861-1868.
4. Warnock G., Rajotte R. Human pancreatic islet transplantation. Transplantation Reviews 1992;6:195-208.
5. Remuzzi F., Ruggenenti P., Mauer S. Pancreas and kidney/pancreas transplants; experimental medicine or real improvement? Lancet 1994;343:27-31.
6. Weber C., Hardy M., Riveria S., Bailey-Braxton D., Michler R., Thomas W., Chabot J., Pi-Sunyer F., Wood M., Reemtsma K. Diabetic mouse bioassay for functional and immunologic human and primate islet xenograft survival. Transplant proceedings 1986;18:823-828.
7. Weber C., Zabinski S., Koschintzky T., Rajotte R., Wicker L., D'Agati V., Peterson L., Norton J., Reemtsma K. The role of CD4⁺ helper T cells in destruction of microencapsulated islet xenografts in NOD mice. Transplantation 1990;49:396-404.

354320 5984050

8. Mandel T., Koulmanda M., Loudovaris R., Bacej A. Islet grafts in NOD mice: A comparison of iso-, allo, and pig xenografts. Transplant Proceedings 1989;21:3813-3814.
9. Weber C., D'Agati V., Ward L., Costanzo M., Rajotte R., Reemtsma K. Humoral reaction to microencapsulated rat, canine, porcine islet xenografts in spontaneously diabetic NOD mice. Transplantation Proceedings 1993;25:462-463.
10. Akita K., Ogawa M., Mandel T. Effect of FK506 and anti-CD4 therapy on fetal pig pancreas xenografts and host lymphoid cells in NOD/Lt, CBA, and BALB/c mice. Cell Transplantation 1994;3:61-73.
11. Morris C., Fung M., Simeonovic S., Wilson D., Hapel A. Cyokine expression in CDA.H mice following xenotransplantation of fetal pig proislets. Transplantation Proceedings 1994;26:1304-1305.
12. Lenschow D., Zeng Y., Thistlewaite J., Montag A., Brady W., Gibon M., Linsley P., Bluestone J. Long-term survival of xenogeneic pancreatic islet grafts induced by CTLAlg. Science 1992;257:789-795.
13. Falqui L., Finke E., Carel J., Scharp D., Lacy P. Marked prolongation of human islet xenograft survival (human-to-mouse) by low temperature culture and temporary immunosuppression with human and mouse antilymphocyte sera. Transplantation 1991;51:1322-1325.
14. Ricordi C., Scharp D., Lacy P. Reversal of diabetes in nude mice after transplantation of fresh and seven

day cultured (24°C) human pancreatic islets.
Transplantation 1988;45:994-996.

15. Ricordi C., Kneteman N., Scharp D., Lacy P.
5 Transplantation of cryopreserved human pancreatic
islets into diabetic nude mice. World J Surgery
1988;12:861-865.
16. Ricordi C., Lacy P., Sterbenz K., Davie J. Low-
10 temperature culture of human islets or in vivo
treatment with L3T4 antibody produces a marked
prolongation is islet (human-to mouse).
Transplantation 1987;44:465-468.
- 15 17. Ricordi C., Lacy P., Sterbenz K., Davie M. Low-
temperature culture of human islets plus in vivo
treatment of L3T4 antibody produces a marked
prongation of islet human-to-mouse xenograft
survival. Proc Natl Acad Sci 1987;84:8080-8084.
- 20 18. Ricordi C., Finke E., Lacy P. A method for the mass
isolation of islets from the adult pig pancreas.
Diabetes 1986;35:649.
- 25 19. Weber C., Costanzo M., Zabinski S., Krekun S.,
Koschitzky T., D'Agati V., Wicker L., Rajotte R.,
Reemtsma K. Xenografts of microencapsulated rat,
canine, porcine, and human islets into streptozotocin
(SZN)-and spontaneously diabetic NOD mice. In:
30 Ricordi C. (Eds.) Pancreatic Islet Transplantation,
R.G. Landes, Austin, 1992:177-190.
20. Weber C., Reemtsma K. Microcapsulation in small
animals-II: Xenografts. In: Lanza R., Chick W.
35 (Eds.). PANCREATIC ISLET TRANSPLANTATION SERIES; VOL

004085 59864060

III: IMMUNOISOLATION OF PANCREATIC ISLETS. R.
Landes, Austin, 1994:59-79.

21. Zhou D., Sun Y., Vacek I., Ma P., Sun A.
5 Normalization of diabetes in cynomolgus monkeys by
xenotransplantation of microencapsulated porcine
islets. Transp Proc 1994;26:1091-1092.
22. Giannarelli R., Marchetti P., Villani G., DiCarlo A.,
10 Cosimi S., Andreozzi M., Cruschelli L., Masioco P.,
Coppelli A., Navalesi R. Preparation of pure, viable
porcine and bovine islets by a simple method.
Transplantation Proceedings 1994;26:630-631.
23. Jos C., Connolly J., Deardon D., Pearson R., Parrot
15 N., Johnson R. A simple method for isolation from
the rabbit pancreas. Transplantation 1994;58:390-
392.
24. Reemtsma K., Weber C., Kazin M., Pi-Sunyer F.,
20 Nilaver G., Fenoglio C. Xenogeneic islets of
Langerhans for human transplantation: Functional and
morphologic studies of primate, bovine and rabbit
islets. In: Friedman E., L'Esperance L. (Eds).
25 Diabetic-Renal-Retinal Syndrome Vol. 3, Grune and
Stratton, New York, 1986:521-546.
25. Forty J., Cary N., White D., Wallwork J. Hyperacute
30 rejection of rabbit hearts by human blood is mediated
by the alternative pathway of complement.
Transplantation Proceedings 1992;24:488-489.
26. Platt J., Back F. The barrier to
35 xenotransplantation. Transplantation 1991;52:937-
947..

27. Auchincloss H. Xenogeneic transplantation.
Transplantation 1988;46:1-20
- 5 28. Lanza R.P., Beyer A.M., Chick W.L. Xenogeneic
humoral responses to islets transplanted in biohybrid
diffusion chambers. Transplantation 1994;57:1371-
1375.
- 10 29. Faustman F., Coe C. Prevention of xenograft
rejection by masking donor HLA class I antigens.
Science 1991;252:1700-1702.
- 15 30. Colton C., Avgoustiniatos E. Bioengineering in
development of the hybrid artificial pancreas. J.
Biochem Enf. 1991;113:152-170.
- 20 31. Lanza R., Sullivan S., Chick W. Islet
transplantation with immunoisolation. Diabetes
1992;41:1503-1510.
- 25 32. Scharp D., Swanson C., Olack B., Latta P., Hegre O.,
Doherty E., Gentile F., Flavin K., Ansara M., Lacy P.
Protection of encapsulated human islets implanted
without immunosuppression in patients with Type I or
Type II diabetes and in nondiabetic control subjects.
Diabetes 1994;43:1167-1170.
- 30 33. Ricker A., Stockberger S., Halban P., Eisenbarth F.,
Bonner-Weir S. Hyperimmune response to
microencapsulated xenogeneic tissue in non obese
diabetic mice. In: Jaworski M. (Eds). The Immunology
of Diabetes Mellitus. Elsevier, Amsterdam, 1986:193-
200.
- 35 34. Weber C., Zabinski S., Norton J., Koschitzky T.,

5

35. Halle J., Bourassa S., Leblond F., Chevalier S., Beaudry M., Chapdelaine A., Cousineau S., Saintonge J., Yale J. Protection of islets of langerhans from antibodies by microencapsulation with alginate-poly-
10 l-lysine membranes. Transplantation 1993;44:350-354.
36. Horcher A., Zekorn T., Siebers U., Klock G., Frank H., Houben R., Bretzel R.G., Zimmerman U., Federlin K. Transplantation of microencapsulated islets in
15 rates: Evidence for induction of fibrotic overgrowth by islets alloantigens released from microcapsules. Transpl Proc 1994;26:784-786.
37. Sibers R., Zekorn T., Horcher A., Klock G., Houben R., Frank H., Bretzel R.G. , Zimmerman U., Federlin K. Microencapsulated transplantation of allogeneic
20 islets into specifically presensitized recipients. Transpl Proc 1994;26:787-788.
38. Lanza R., Kuhtreiber W., Ecker D., Staruk J., Chich W., Xenotransplantation of porcine and bovine islets without immunosuppression using uncreated alginate
25 microspheres. Transplantation 1995;59:1377-1384.
39. Lanza R., Ecker D., Kuhtrieber W., Staruk J., Marsh J., Chick W. A simple method for transplanting discordant islets into rats using alginate gel
30 spheres. Transplantation 1995;59:1485-1487.
40. Soon-Shiong P., Feldman E., Nelson R., Komtebedde J.,

Smidsrod O., Skauk-Braek G., Espevik T., Heintz R.,
Lee M. Successful reversal of spontaneous diabetes
in dogs by using intraperitoneal microencapsulated
islets. Transplantation 1992;54:769-774.

5

41. Soon-Shiong P., Heintz R.E., Merideth N., Yao Q.X.,
Yao Z., Zheng T., Murphy M., Moloney M.K., Schmehl
M., Harris M., Mendez R., Sandford P.A. Insulin
independence in a type I diabetic patient after
encapsulated islet transplantation. Lancet
1994;343:950-951.

10

42. Heald K., Jay T., Downing R. Assessment of the
reproducibility of alginate encapsulation of
pancreatic islets using the MTT colorimetric assay.
Cell transplantation 1994;3:333-337.

15

43. Jarpe A., Hickman M., Anderson J. Flow cytometric
enumeration of mononuclear cell populations
infiltrating the islets of Langerhans in prediabetic
NOD mice: Development of model of autoimmune
insulinitis for Type I diabetes. Regional Immunology
1990;3:305-317.

20

44. Miller B., Appal M., O'Neil J., Wicker L. Both the
Lyt-2+ and L3T4+T cell subsets are required for the
transfer of diabetes in nonobese diabetic mice. J.
Immunol 1988;140:52-58.

25

45. Haskins K., Portas M., Bradley B. T-lymphocyte clone
specific for pancreatic islet antigen. Diabetes
1988;37:1444-1448.

30

46. Jansen A., Homo-Delarche F., Hooijkaas J., Leenen P.,
Dardenne M., Drexhage H. Immunohistochemical

35

00220-5964060

characterization of monocytes-macrophages and dendritic cells involved in the initiation of the insulinitis and B-cell destruction in NOD mice. Diabetes 1994;43:667-674.

5

47. Prins J., Todd J., Rodrigues N., Ghosh S., Hogarth M., Wicker L. Gaffney E., Fishcer P., Sirotina A., Peterson L. Linkage on Chromosome 3 of autoimmune diabetes and defective Fc receptor for IgG in NOD mice. Science 1993;260:695-695.
48. Lipes M., Rosenzweig A., Tan K., Tanigawa G., Lass D., Seidman J., Eisenbarth G. Progression to diabetes in nonobese diabetic (NOD) mice with transgenic T cell response. Science 1993;259:1165-1169.
49. Gelber C., Pabrosky L., Singer S., McAteer D., Tisch R., Jolicoeur C., Buelow R., McDevitt H., Fathman G. Isolation of nonobese diabetic mouse T-cells that recognize novel autoantigens involved in the early events of diabetes. Diabetes 1994;43:33-39.
50. Podolin P., Pressey A., DeLarato N., Fischer P., Peterson L., Wicker L. I-E+ nonobese diabetic mice develop insulinitis and diabetes. J Exp Med 1993;178:793-803.
51. Haskins K., Portas M., Bergman B., Lafferty K., Bradley B. Pancreatic islet-specific T-cell clones from nonobese diabetic mice. PNAS 1989;86:8000-8004.
52. Haskins K., McDuffie M. Acceleration of diabetes in young NOD mice with a CD4⁺ islet-specific T cell clone. Science 1990;249:1433-1436.

53. Supon P., Stecha P., Haskins K. Anti-islet cell antibodies from NOD mice. *Diabetes* 1990;39:1366-1392.
- 5 54. Bergman B., Haskins K. Islet-specific T-cell clones from the NOD mouse respond to B-granule antigen. *Diabetes* 1994;43:197-203.
- 10 55. Peterson J., Pike B., McDuffie M., Haskins K. Islet-specific T-cell clones transfer diabetes to nonobese diabetic (NOD) F₁ mice. *J. Immunol* 1994;153:2800-2806.
- 15 56. Kaufman D., Clare-Salzier M., Tian J. Spontaneous loss of T-cell tolerance to glutamic acid decarboxylase in murine insulin-dependent diabetes. *Nature* 1993;365:69-72.
- 20 57. Tisch R., Yang X., Singer S., Liblau R., Fugger L., McDevitt H. Immune response to glutamic acid decarboxylase correlates with insulinitis in non-obese diabetic mice. *Nature* 1993;366:72-75.
- 25 58. Dylan D., Gill R., Schloot N., Wegmann D. Epitope specificity, cytokine production profile and diabetogenic activity so insulin-specific T cell clones isolated from H=NOD mice. *Eur J. Immunology* 1995;25:1062.
- 30 59. Wang Y., Hao L., Gill R., Lafferty K. Autoimmune diabetes in NOD mouse is L3T4 T-lymphocyte dependent. *Diabetes* 1987;36:535-538.
- 35 60. Boitard D., Bendelac A., Richard M., Carnaud C., Bach J. Prevention of diabetes in nonobese diabetic mice

004986-03798

5 61. Serrese D., Gaskins H., Leiter E. Defects in the
differentiation and function of antigen presenting
cells in NOD/lt. Mice. J Immunol 1993;150:2534-2543.

10 62. Rabinovitch A., Sumoski W., Rajotte R., Warnock G.
Cytotoxic effects of cytokines on human pancreatic
islet cells in monolayer culture. J. Of Clinical
Endocrinology and Metabolism 1990;71:152-156.

15 63. Bergmann L., Kroncke K., Suschek D., Kolb H., Kolb-
Bachofen C. Cytotoxic action of IL-1B against
pancreatic islets in mediated via nitric oxide
formation and is inhibited by N^G -monomethyl-L-
arginine. FEBS letter 1992;299:103-106.

20 64. Xenos D., Stevens R., Gores P., Casanova D., Farney
A., Sutherland D., Platt J. IL-1 induced inhibition
of B-cell function is mediated through nitric oxide.
Transpl Proc 1993;25:994-994.

25 65. Anderson H., Jorgensen K., Egeberg J. Mandrup-Poulsen
R., Berup J. Nicotinamide prevents Interleukin-1
effects on accumulated insulin release and nitric
oxide production in rat islets of Langerhans.
Diabetes 1994;43:770-777.

30 66. Christianson S., Shultz L., Leiter D. Adoptive
transfer of diabetes into immunodeficient NOD-
Scid/Scid mice. Diabetes 1993;42:44-45.

35 67. Gerling I., Friedman H., Greiner D., Schultz L.,

Leither E. Multiple low-dose streptozotocin-induced diabetes in NOD-Scid/Scid mice in the absence of functional lymphocytes. Diabetes 1994;43:433-440.

- 5 68. Rohane P., Shimada A., Kim D., Edwards C., Charlton B., Shultz L., Fathman C. Islet infiltrating lymphocytes from prediabetic NOD mice rapidly transfer diabetes to NOD-scid/scid mice. Diabetes 1995;44:550-554.

10

69. Shultz L., Schweitzer P., Christian S., Gott B., Schweitzer I., Tennent B., McKenna S., Mobraaten L., Rajan R., Greiner D., Leiter E. Multiple defects in innate and adaptive immunologic function in NOD/LtSz-scid mice. J of Immunology 1995;154:180-191.

15

70. Weber C., Krekun S., Loschitzky S., Zabinski S., D'Agati C., Hardy M., Reemtsma K. Prolonged functional survival of rat-to-NOD mouse islet xenografts by ultraviolet-B (UV-B) irradiation plus microencapsulation of donor islets. Transplantation Proceedings 1991;23:764-766.

20

71. Lafferty K. Circumventing rejection of islet grafts; An overview. In: Van Schifgaarde R., Hardy M. (Eds). Elsevier, Amsterdam, 1988:279-291.

25

72. Ricker A., Bhatia V., Bonner-Weir S., Eisenbarth G. Microencapsulated xenogeneic islet grafts in NOD mouse: Dexamethasone and inflammatory response. Diabetologia 1989;32:53.

30

73. Lafferty K.J., Hao L. Approaches to the prevention of immune destruction of transplanted pancreatic islets. Transpl Proc 1994;26:399-400.

35

0004955-0329

74. Weber C., Price J., Costanzo M., Becker A., Stall A.
NOD mouse peritoneal cellular response to poly-L-lysine-alginate microencapsulated rat islets.
Transplantation Proceedings 1994;26:1116-1119.
- 5
75. Weber C., Tanna A., Costanzo M., Price J., Peterson L., Wicker L. Effects of host genetic background on survival of rat → mouse islet xenografts.
Transplantation Proceedings 1994;26:1186-1188.
- 10
76. Wijsman J., Atkinson P., Mazheri R. Histological and immunopathological analysis of recovered encapsulated allogeneic islets from transplanted diabetic BB/W rates. Transplantation 1992;54:588-592.
- 15
77. Cole D., Waterfall M., McIntyre M., Baird J. Microencapsulated islet grafts in the BB.E rat a possible role for cytokines in graft failure. Diabetologia 1992;35:231-237.
- 20
78. Mazaheri R., Atkinson P., Stiller C., Dupre J., Vose J., O'Shea F. Transplantation of encapsulated allogeneic islets into diabetic BB/W rates: Effects of immunosuppression. Transplantation 1991;51:750-754.
- 25
79. Darquy S., Reach G. Immunoisolation of pancreatic B cells by microencapsulation. Diabetologia 1985;28:776-780-.
- 30
80. Baldwin W., Pruitt S., Brauer R., Daha M., Sanfillippo F. Complement in organ transplantation. Transplantation 1995;59:797-808.
- 35
81. Iwata H., Takagi R., Amemiya H. Marked prolongation of islet xenograft survival (hamster-to-mouse) by

0004935-0327060

microencapsulation and administration of 15-deoxyspergualin. Transplantation Proceedings 1992;24:1516-1518.

- 5 82. Pierson R., Winn H., Russell P., Auchincloss H. CD-4 positive lymphocytes play a dominant role in murine xenogeneic responses. Transplantation Proceedings 1989;21:519-521.
- 10 83. Gill R., Wolf L., Daniel D., Coulombe M. CD4+ T cells are both necessary and sufficient for islet xenograft rejections. Transp Proc 1994;26:1203-1204.
- 15 84. Weber C., Zabinski S., Koschitzky T., Wicker L., Rajotte R., Peterson L., D'Agati V., Reemtsma K. Microencapsulated dog and rat islet xenografts into streptozotocin-diabetic and NOD mice. Hoem Metab Res 1990;35:219-226.
- 20 85. Cirulli V., Halban P., Rouiller D. Tumor necrosis factor a modifies adhesion properties of rat islet B cells. J Clin Invest 1993;92:1868-1876.
- 25 86. Campbell I., Iscara A., Harrison L. Interferon gamma and tumor necrosis factor alpha: cytotoxicity to murine islets of Langerhans. J Immunol 1988;141:1325-1329.
- 30 87. Rabionovitch A. Immunoregulatory and cytokine imbalances in the pathogenesis of IDDM. Diabetes 1994;43:613-621.
- 35 88. Ablamunits V., Baranove F., Mandrup-Poulsen R., Nerup J. In vitro inhibition of insulin release by blood mononuclear cells from insulin-dependent diabetic and healthy subjects: synergistic action of IL-1 and TNF.

00049865-032798

Cell Transplantation 1994;3:55-60.

89. Mandrup-Poulsen T., Bendtzen D., Dinarello C., Nerup J. Human tumor necrosis factor potentiates interleukinmediated rate of pancreatic B-cell cytotoxicity. J Immunol 1987;139:4077-4082.
90. Thai N., Wang S., Valdivia L., Celli S., Reilly M. Demetris A., Simmons R., Stazi T., Fung J. Cytokine messenger RNA profiles in hamster-to-rat liver xenografts. Transplantation Proceedings 1993;25:444-445.
91. O'Connell P., Pacheco-Silca A., Nickerson P., Muggia R., Bastos M., Kelly V., Strom R. Unmodified pancreatic islet allograft rejection results in preferential expression of certain T cell activation transcripts. J Immunol 1993;150:1093-1104.
92. Nickerson P., Pacheco-Silca A., O'Connell P., Steurer W., Kelly V., Strom R. Analysis of cytokine transcripts in pancreatic islet cell allografts during rejection and tolerance induction. Transplantation Proceedings 1993;25:984-985.
93. Lowry R., Takeuchi T. The Th1, Th2 paradigm and transplantation tolerance R. Landes , Austin. 1994; In press.
94. Lederer J., Liou J., Todd M., Glicher L., Lichtman A. Regulation of cytokine gene expression in T helper cell subsets. J Immunol 1994;1:78-86.
95. Janeway C., Bottomly K. Signals and signs for lymphocyte responses. Cell 1994;76:275-285.

96. Aguilar-Diosdada M., Parkinson D., Corbett J., Kwon G., Marshall C., Gingerich R., Santiafo J., McDaniel M. Potential autoantigens in IDDM: expression of carboxypeptidase-H and insulin but not glutamate decarboxylase on the B-cell surface. Diabetes 1994;43:418-425.
97. Jenkins M. The ups and downs of T cell costimulation. Immunity 1994;4:443-446.
98. Liu Y., Jones B., Brady W., Janeway C., Linsley P. Co-stimulation of murine CD4 T cell growth: cooperation between B7 and heat-stable antigen. Eur J Immunol 1992;22:2855-2859.
99. Harding F., McArthur J., Fross J., Raulet D., Alliston J. CD28-mediated signaling co-stimulates murine T cells and prevents induction of anergy in T-cell clones. Nature 1992;356:607-609.
100. Galvin F., Greeman G., Razi-Wolf Z., Hall W., Benacerraf B., Nadler L., Reiser H. Murine B7 antigen provides a sufficient costimulatory signal for antigen-specific and MHC-restricted T cell activation. J Immunol 1992;149:3802-3808.
101. Hathcock K., Laszio G., Dickler H., Bradshaw J., Linsley P., Hodes R. Identification of an alternative CTLA-4 ligand costimulatory for T cell activation. Science 1993;262:905-907.
102. Perrin P., Scott D., Quigley L., Albert P., Feder O., Gray G., Abe R., June C., Racke M. Role of B7: CD28/CTLA-4 in the induction of chronic relapsing experimental allergic encephalomyelitis. J of

Table 1. <i>Continued</i>					
Variable	Mean	SD	Median	Mode	Range
Age (years)	45.2	12.5	42.0	40.0	20-75
Gender					
Male	55.0	5.0	55.0	55.0	40-70
Female	45.0	5.0	45.0	45.0	30-60
Education (years)	12.5	2.0	12.0	12.0	8-16
Occupation					
Professional	30.0	10.0	30.0	30.0	20-50
Managerial	20.0	10.0	20.0	20.0	10-40
Technical	10.0	10.0	10.0	10.0	0-30
Service	10.0	10.0	10.0	10.0	0-30
Unemployed	10.0	10.0	10.0	10.0	0-30
Retired	10.0	10.0	10.0	10.0	0-30
Other	10.0	10.0	10.0	10.0	0-30
Marital status					
Married	60.0	10.0	60.0	60.0	40-80
Single	20.0	10.0	20.0	20.0	10-40
Divorced	10.0	10.0	10.0	10.0	0-30
Widowed	10.0	10.0	10.0	10.0	0-30
Other	10.0	10.0	10.0	10.0	0-30
Religion					
Christian	70.0	10.0	70.0	70.0	50-90
Muslim	20.0	10.0	20.0	20.0	10-40
Hindu	10.0	10.0	10.0	10.0	0-30
Jewish	10.0	10.0	10.0	10.0	0-30
Other	10.0	10.0	10.0	10.0	0-30
Income (USD/month)	1000.0	500.0	1000.0	1000.0	500-2000
Health status					
Good	60.0	10.0	60.0	60.0	40-80
Fair	20.0	10.0	20.0	20.0	10-40
Poor	10.0	10.0	10.0	10.0	0-30
Very poor	10.0	10.0	10.0	10.0	0-30
Other	10.0	10.0	10.0	10.0	0-30

- | Table 1. <i>Continued</i> | | | | | |
|---------------------------|------|------|--------|------|-------|
| Variable | Mean | SD | Median | Mode | Range |
| Age (years) | 45.2 | 12.5 | 42.0 | 40.0 | 20-75 |
| Gender | | | | | |
| Male | 55.0 | 5.0 | 55.0 | 55.0 | 40-70 |
| Female | 45.0 | 5.0 | 45.0 | 45.0 | 30-60 |
| Education (years) | 12.5 | 2.0 | 12.0 | 12.0 | 8-16 |
| Occupation | | | | | |
| Professional | 30.0 | 10.0 | 30.0 | 30.0 | 20-50 |
| Managerial | 20.0 | 10.0 | 20.0 | 20.0 | 10-40 |
| Technical | 10.0 | 10.0 | 10.0 | 10.0 | 0-30 |
| Service | 20.0 | 10.0 | 20.0 | 20.0 | 10-40 |
| Unemployed | 10.0 | 10.0 | 10.0 | 10.0 | 0-30 |
| Marital status | | | | | |
| Married | 60.0 | 10.0 | 60.0 | 60.0 | 40-80 |
| Single | 20.0 | 10.0 | 20.0 | 20.0 | 10-40 |
| Divorced | 10.0 | 10.0 | 10.0 | 10.0 | 0-30 |
| Widowed | 10.0 | 10.0 | 10.0 | 10.0 | 0-30 |
| Religion | | | | | |
| Christian | 70.0 | 10.0 | 70.0 | 70.0 | 50-90 |
| Muslim | 20.0 | 10.0 | 20.0 | 20.0 | 10-40 |
| Hindu | 10.0 | 10.0 | 10.0 | 10.0 | 0-30 |
| Other | 0.0 | 0.0 | 0.0 | 0.0 | 0-1 |

between CD40 and its ligand gp39 in the development of murine lupus nephritis. The American Association of Immunologists 1995;154: 1470-1480.

- 5 111. Gin H., Cadic C., Baquey C., Dupuy B. Peritoneal exudates from microencapsulated rat islets of Langerhans xenografted mice presenting characteristics of potentially cytotoxic non-specific inflammation. J of Microencapsulation 1992;9: 489-494.
- 10 112. Clayton H., London N., Colloby P., Bell P., James R. The effect of capsule composition on the biocompatibility of alginate-poly-l-lysine capsules. J. Microencapsulation 1991;8: 221-233.
- 15 113. Soon-Shioh P., Oterlie M., Skjak-Braek G., Smidsrod O., Heintz R., Lanza R.P., Espevik T. An immunologic basis for the fibrotic reaction to implanted microcapsules. Transpl Proc 1991;23: 758-759.
- 20 114. DeVos P., Wolters G., VanSchilfgaarde R. Possible relationship between fibrotic overgrowth of alginate-polysine-alginate microencapsulated pancreatic islets and the microcapsule integrity. Transp Proc 1994;26: 782-783.
- 25 115. Chang T. Artificial cells in immobilization biotechnology. Art Cells & Immob Biotech 1992;20: 1121-1143.
- 30 116. Lum Z., Tai I., Krestow M., Norton J., Vacek I., Sun A. Prolonged reversal of diabetic state in NOD mice by xenografts of microencapsulated rat islets. Diabetes 1991;40: 1511-1516.
- 35

117. Chicheportiche D., Reach G. In vitro kinetics of insulin release by microencapsulated rat islets: effects of the size of the microcapsules. *Diabetologia* 1988;31: 54-57.
- 5
118. Weber C., Constanzo M., Kredun S., D'Agati V. Causes of destruction of microencapsulated islet grafts: Characteristics of a 'double-wall' poly-l-lysine-alginate microcapsule. *Diabetes, Nutrition and Metabolism* 1993;1: 167-171.
- 10
119. Vanenbossche G., Van Oostveldt P., Demeester J., Remon J. The molecular weight cut-off of microcapsules is determined by the reaction between alginate and polylysine. *Biotechnology and Bioengineering* 1993;42: 381-386.
- 15
120. Linderman G., Adams J., Cory S., Harris A. B-lymphoid to granulocytic with during hematopoiesis in a transgenic mouse strain. *Immunity* 1994;1: 517-527.
- 20
121. Takeuchi T., Lowry R., Konoieczny B. Heart allografts in murine systems. *Transplantation* 1992;53: 1281-1294.
- 25
122. Ravetch J., Kinet J. *J Ann Rev Immunol* 1993;9: 457-492.
123. Takai T., Li M., Sylvestre D., Clynes R., Ravetch J. FcR γ chain deletion results in pleiotropic effector cell defects. *Cell* 1994;76: 519-529.
- 30
124. Krych M., Atkinson J., Holers v. Complement receptors. *Curr Opin Immunol* 1992;4: 8-13.
- 35

004985 032799
00220 59864060

125. Frank M., Fries: The role of complement in inflammation and phagocytosis. *Immunol Today* 1991;12: 322-326.
- 5 126. Whiteley P.J., Jensen P.E. Pierce C.W., Abruzzini A.F., Kapp J.A. Helper T-cell clones that recognize autologous insulin are stimulated in nonresponder mice by pork insulin. *Proc Natl Acad Sci USA* 1998;85: 2723-2727.
- 10 127. Jensen P.E., Kapp J.A. Stimulation of helper T cells and dominant suppressor T cells that recognize autologous insulin. *J Mol Cell Immunol* 1985;2: 133-133.
- 15 128. Poindexter N.J., Landon C., Whiteley P.J., Kapp J.A. Comparison of the T cells receptors on insulin-specific hybridomas from insulin transgenic and nontransgenic mice. Loss of a subpopulation of self-reactive clones. *J Immunol* 1992;149: 38-44.
- 20 129. Ke Y., Li Y., Kapp J.A. Ovalbumin injected with complete Freund's adjuvant stimulates cytolytic responses. *Eur J Immunol* 1995;25: 549-553.
- 25 130. Seder R., Paul W. Acquisition of lymphokine-producing phenotype by CD 4⁺ T cells. *Annu Rev Immunology* 1994;12: 635-635.
- 30 131. Beckerman K., Rogers H., Corbett J., Schreiber R., McDaniel M., Unanue E. Release of nitric oxide during the T-cell independent pathway of macrophage activation *J Immunol* 1993;150: 888-895.
- 35 132. Weinberg B., Granger D., Pisetsky D., Seldin M.,

054300 59364060

- Misukonis M., Mason N., Pippen A., Ruiz P., Wood E., Gilkeson G. The role of nitric oxide in the pathogenesis of spontaneous murine autoimmune disease: Increased nitric oxide production and nitric oxide synthase expression in MRL-lpr/lpr mice, and reduction of spontaneous glomerulonephritis and arthritis by orally administered N^g-monomethyl-L-arginine. J. Exp Med 1994;179: 651-660.
133. Ravetch J. Fc Receptors: Rubor Redux. Cell 1994;78: 553-560.
134. Unreels J., Scigliano E., Freedman V. Structure and function of human and murine receptors for IgG. Annual Review of Immunology 1988;6: 251-281.
135. Platt J., Lindman B., Geller R., Noreen H., Swanson J., Dalmaso A., Bach F. The role of natural antibodies in the activation of xenogenic endothelial cells. Transplantation 1991;52: 1027-1043.
136. Baxter G., Cooke A. Complement lytic activity has no role in the pathogenesis of autoimmune diabetes in NOD mice. Diabetes 1994;42: 1574-1578.
137. Moses R., Winn H., Auchincloss H. Evidence that multiple defects in cell-surface molecule interactions across species differences are responsible for diminished xenogenic T cell responses. Transplantation 1992;53: 203-209.
138. Moses R., Pierson R., Winn H., Auchincloss H. Xenogeneic proliferation and lymphokine production are dependent on CD4+ helper T cells and self antigen-presenting cells in the mouse. J Exp Med 1990;172:

567-575.

139. Hardy M., Lau H., Weber C., Reemtsma K. Pancreatic islet Transplantation; Induction of graft acceptance by ultraviolet irradiation of donor tissue. Ann Surg 1984;200: 441-450.
140. Sullivan F., Ricordi C., Hauptfeld V., Lacy P. Effect of low-temperature culture and site of transplantation on hamster islet xenografts survival (hamster-to-mouse). Transplantation 1987;44: 465-468.
141. Hering B., Bretzel R., Federlin K., Horm Metabol Res 1988;20:537-545.
142. Lacy P., Lacy E., Finke E., Yasunami Y. Diabetes 1982;31:109-111.
143. Nathan D. NEJM 1993;328:176-1685.
144. Robertson R. NEJM 1992;327: 1861-1868.
145. Linsley P.S., Brady W., Umes M., Grosmaire L.S., Damie N.K., Ledbetter J.A. CTLA-4 is a second receptor for the B cell activation antigen B7. J Exp Med 1991; 74: 561-569.
146. Korbitt G.S., Ao Z., Warnock G.L., Flashner M., Rajotte R.V. Successful reversal of diabetes in nude mice by transplantation of microencapsulated porcine neonatal islet cell aggregats. Transplantation Proceedings 1995;27:3212.
147. Korbitt G.S., Ao Z., Warnock G.L., Rajotte R.V. Large-scale isolation of viable porcine neonatal islet cell

8622E0 59864060

Second Series of Experiments

The goal of this study was to clarify the mechanism(s) of destruction of microencapsulated islet xenografts by spontaneously diabetic NOD mice, the best available model of human insulin-dependent diabetes (IDDM). We have found that NOD helper T-cells and MHC both are necessary for destruction of encapsulated islets (30,31), and we have documented that empty microcapsules are biocompatible in NOD mice (27-29,31). Encapsulated islet xenografts biopsied at rejection in NOD mice contained abundant peri-capsular macrophages and immunoglobulins, with IL- 1, $\text{TNF}\alpha$, and IL-4 messenger RNA (13,28,29). Therefore, we postulated that NOD rejection was initiated by donor antigens that were secreted from encapsulated islets, and were processed via the MHC class II pathway by host APC. NOD CD4+ T-cells then promoted a Th2 response, with donor islet destruction occurring via cytokine-mediated events.

Involvement of APC in immune responses to islet xenografts was suggested by recent studies of Lenschow et al. (15), who found that blockade of the costimulatory molecule, B7 with the soluble fusion protein, CTLA4-Ig, prolonged human-to-mouse islet xenografts in SZN-diabetic mice. Several studies, in vitro and in vivo, have shown that foreign molecules that interact with the T-cell fall on their own to stimulate naive T-cells to proliferate and may induce antigen-specific anergy. At least one additional (costimulatory) signal is required, and it is delivered by APC. In mice, one such costimulatory pathway involves the interaction of the T-cell surface antigen, CD28 with either one of two ligands, B7-1 and B7-2, on APCs (4,7,8,10,11,17,22). Once this full interaction of T-cells and APC occurs, reexposure of T-cells to the peptide, mitogen, etc., will result in proliferation

CTLA-4 is a cell surface protein similar to CD28; however, unlike CD28, CTLA-4 is expressed only on activated T-cells. B7-1 has a higher affinity for CLTA4 than CD28, and it has been suggested that CTLA4 may modulate functions of CD28 (5,6,11). CTLA4-Ig is a recombinant soluble fusion protein, with the extracellular binding domain of the CTLA4 molecule and the constant region of the IgG1 gene, which inhibits T-lymphocyte responses in mice (9,14). Administration of CTLA4-Ig to mice induces antigen-specific unresponsiveness (7,11,25), and long-term acceptance of murine cardiac allografts (3,21). In addition, CTLA4-Ig has been reported to reduce the incidence of diabetes in NOD mice (16). We have recently found that murine CTLA4-Ig prolonged survival of encapsulated adult rabbit islets in NOD mice (26).

Neonatal porcine islets were isolated from White Landrace pigs and tissue cultured as previously described (12). Approximately 8000 islets were encapsulated in double-wall, Poly-L-lysine-alginate microcapsules and grafted intraperitoneally in NOD or NOD-SCID mice, as previously reported (29,31). Controls received approximately 8000 unencapsulated neonatal islets grafted beneath the splenic or renal capsule.

30 Murine CLTA4-Ig, provided by Bristol-Myers-Squibb (P.S.L.) Seattle, WA, was administered at 200 μ g intraperitoneally (i.p.), day zero and then q.o.d. for 21 days, or until graft rejection if that occurred prior to
35 day #21. Graft function was monitored daily by measurement of random blood glucose for 2 wk and then weekly (31). Graft rejection was defined as random blood

glucose >250 mg/dL for 2 consecutive days.

Biopsies of long-term functioning peritoneal micro-
capsules were done periodically, using metaflane anes-
5 thesia and a sterile technique. Removal of 100-200
microcapsules allowed histologic light microscopic and
insulin immunochemical studies without altering graft--
related normoglycemia (31). Statistical differences
between groups were assessed by use of Student's t-test
10 and by ANOVA.

RESULTS

Neonatal pig islets are actually dispersed neonatal
15 porcine pancreatic cells that reaggregate in tissue
culture to form "islet"-like spheroids with
approximately 5-10% beta cells (see Fig. 16), which is
significantly higher than the 1-2% beta cell
concentration in the adult porcine pancreas.
20 Approximately 30,000-100,000 islets were obtained from
each 2-5-day-old neonatal donor pig (see Fig. 14).
Neonatal pig islet cells secreted insulin in vitro
after microencapsulation (see Fig. 15). Biopsies of
these "islets" >100 days following transplantation to
25 streptozotocin-diabetic NOD-Scid mice revealed increased
numbers of intensely insulin-positive islet cells in
most cell aggregates (Fig. 28). Thus, neonatal pig
islets appear to differentiate and divide during the
transplant period.

30 Treatment of NOD mice with CTLA4-Ig significantly
prolonged survival of intraperitoneal poly-L-lysine-
alginate microencapsulated donor neonatal porcine islet
xenografts (CAP/I.P.) in spontaneously diabetic NOD
35 mice, when compared to either islet microencapsulation
or NOD CTLA4-Ig treatment alone (Table 7). Biopsies of
long-term functioning encapsulated neonatal porcine

islets from CTLA4-Ig-treated NOD mice documented intact microcapsules, containing viable donor islets, with many insulin-positive beta cells, and no peri-capsular NOD cellular response (Fig. 29). Biopsies of NODs controls
5 receiving intraperitoneal encapsulated porcine islets, without CTLA4-Ig treatment uniformly showed pericapsular accumulations of macrophages, neutrophils, and lymphocytes, as previously described (2,20,24). Biopsies of pancreases from NOD mice in all experimental groups
10 showed absence of beta cells, and occasional accumulations of lymphocytes in perivascular areas.

00403E 59864060

- 90 -

Table 7

Survival of microencapsulated (MC) neonatal porcine islets in NOD mice: Effects of NOD treatment with CTLA4-Ig

		Mean Graft survival (days)		
	Technique	Rx.	(n)	x ± SE
Pig-NOD	MC/IP	None	8	27±13
				9, 10, 12, 12, 14, 14, 23, 118 ^(s)
Pig-NOD	MC/IP	CTLA4-Ig	6	110±14*
				74 ^(s) , 80, 101 ^(s) , 108, 137 ^(s) , 161 ^(s)
Pig-NOD	Splenic	CTLA4-Ig	3	5±1
				4, 5, 5
Pig-NOD	Splenic	None	3	6
				5, 6, 7

i.p. = intraperitoneal; ^(s) = sacrifice for biopsy;

*p<0.001 vs. MC alone; CTLA4-Ig, 200 µg i.p., q.o.d. x 3 wk.

DISCUSSION

364220"59864050

The most important finding of this invention is the synergy of donor islet microencapsulation and NOD CTLA4-Ig treatment in prolonging neonatal porcine islet xenograft survival. Neither CTLA4-Ig, nor encapsulation alone was effective. There is evidence that xenorecognition (unlike allorecognition) occurs primarily via the so-called "indirect" antigen presentation pathway, by which host APC present peptides scavenged from extracellular (donor) proteins to host helper T-cells (1,18,19,23). Applicants' recent report, that the host MHC is critical to NOD rejection of encapsulated islet xenografts (30), and applicants' prior observations, that helper NOD T-cells are essential for this response (31), both are consistent with an "indirect" pathway. Applicants' current data suggest that "indirect" antigen presentation may be blocked by CTLA4-Ig in this model of encapsulated islet xenotransplantation.

- 92 -

Table 8

Synergy of Neonatal Porcine Islet Microencapsulation and Xenogeneic NOD Mouse Recipient Treatment with CTLA4Ig

Technique	Rx	(N)	Graft Survival (Days) ($\bar{x} \pm SE$)
MC/IP	None	8	9, 10, 12, 12, 14, 18, 23, 116 (27 \pm 36)
MC/IP	CTLA4Ig	10	61, 74 ^s , 80 ^s , 80, 85, 101 ^s , 108, 137 ^s , 160 ^s , 266 ^s (115.2 \pm 19.3) **
MC/IP	CTLA4Ig*	10	12, 32, 55, 63, 75, 78, 83, 103, 239 ^s , 287 ^s (102.7 \pm 28.2) **
Splenic	CTLA4Ig	3	4, 5, 5 (5 \pm 1)
Splenic	None	3	5, 6, 7 (6 \pm 1)

(s) = sacrifice; **p<.003 vs. MC/IP alone;
MC/IP = microencapsulation/intraperitoneal;
CTLA4Ig* = mutant CTLA4Ig, which does not bind complement.

Pig C-Peptide (ng/ml)

Conclusions

Long-term effect was found with only 21 days of CTLA4Ig. Both wild-type CTLA4Ig and mutant CTLA4Ig* (Y100 F) (which does not bind complement) were effective. (see Figs. 34-35) There was no toxicity to recipients. There is biopsy proof of long-term graft function. (see Figs 30-33) Further proof that long-term grafts are functioning is provided by use of a radio-immunoassay to measure pig insulin (pig C-peptide) as secreted (see Table 9) which is rapidly degraded. Insulin is released by insulin cells that are specific for the pig (pig insulin is detected by the presence of the C-peptide tails). The presence of C-peptide tails in the grafts indicates that the graft is alive and functioning, as exemplified by a long-term graft of 239 days. (see also Figs. 30-31)

In conclusion, applicants have found that neither microencapsulation nor CTLA4-Ig alone prevent NOD destruction of neonatal porcine islets. However, applicants observed synergy between CTLA4-Ig treatment

of NOD recipients and islet encapsulation, with significantly prolonged discordant islet xenograft survival. (see Tables 7 and 8) Because of the availability of large quantities of porcine islets and bioacceptability of the microcapsules and CTLA4-Ig this approach may be clinically relevant (in humans).

03049865 032798

References for the Second Series of Experiments

1. Auchincloss, H. Transplantation 46:1-20; 1988.
2. Barker, C.; Najj, A. N. Engl. J. Med. 327:1861-1868; 1992.
3. Bolling, S.; Turka, L.; Wei, R.; Linsley, P.; Thompson, C.; Lin, H. Transplantation xx:22413-415; 1994.
4. Galvin, F.; Freeman, G.; Razi-Wolf, Z.; et al. J. Immunol. 149:3802-3908; 1992.
5. Gimmi, C.; Freeman, G.; Gribben, J.; Gray, G.; Nadler, L. Immunology 90:6586-6590; 1993.
6. Guerder, S.; Meyerhoff, J.; Flavell, R. Immunity 1:155-166; 1994.
7. Harding, F.; McArthur, J.; Gross, J.; Raulet, D.; Allison, J. Nature 356:607-609; 1992.
8. Hathcock, K.; Laszio, G.; Dickler, H.; Bradshaw, J.; Linsley, P.; Hodes, R. Science 262:905-907; 1993.
9. Hering, B.; Bretzel, R.; Federlin, K. Horm. Metabol. Res. 20:537-545; 1988,
10. Janeway, C.; Bottomly, K. Cell 76:275-285; 1994.
11. Jenkins, M. Immunity 1:443-446; 1994.
12. Korbitt, G.S.; Ao, Z.; Warnock, G.L.; Rajotte, R.V. Transplant. Proc. 27:3267; 1995.

052220"99864060

13. Krych, M.; Atkinson, J.; Holers, V. Curr. Opin. Immunol. 4:8-13; 1992.
14. Lacy, P.; Lacy, E.; Finke, E.; Yasunami, Y. Diabetes 31:109-111, 1982.
15. Lenschow, D.; Zeng, Y.; Thistlethwaite, J.; et al. Science 257:789-795; 1992.
16. Lenschow, D.; Ho, S.; Sattar, H.; et al. J. Exp. Med. 181:1145-1155; 1995.
17. Liu, Y.; Jones, B.; Brady, W.; Janeway, C.; Linsley, P. Eur.J. Immunol. 22:2855-1859; 1992.
18. Moses, R.; Pierson, R.; Winn, H.; Auchincloss, H. J. Exp. Med. 172:567-575; 1990.
19. Moses, R.; Winn, H.; Auchincloss, H. Transplantation 53:203-209; 1992.
20. Nathan, D. N. Engl. J. Med. 328:1676-1685; 1993.
21. Pearson, T.; Alexander, D.; Winn, K.; Linsley, P.; Lowry, R.; Larsen, C. Transplantation 57:1701-1706; 1994
22. Perrin, P.; Scott, D.; Quigley, L.; et al. J. Immunol. 154:1481-1490; 1995.
23. Platt, J., Back, F. Transplantation 52:937-947; 1991.
24. Robertson, R. N. Engl. J. Med. 327:1861-1868; 1992.
25. Satyaraj, E.; Rath, S.; Bal, V. J. Immunol. 155:4669-4675; 1995.

852250"59864060

26. Weber, C.I.; Hagler, M.K.; Chryssochoos, J.T.; et al. Transplant. Proc. 28:821-823; 1996.
27. Weber, C.; Krekun, S.; Koschitzky, S.; et al. Transplant Proc. 23:764-766; 1991.
28. Weber, C.; Price, J.; Costanzo, M.; Becker, A.; Stall, A. Transplant. Proc. 26:1116-1119; 1994.
29. Weber, C.; Reemtsma, K.; Lanza, R.; Chick, W., eds. Austin: R. Landes; 1994:59-79.
30. Weber, C.; Tanna, A.; Costanzo, M.; Price, J.; Peterson, L.; Wicker, L. Transplant. Proc. xxx.???
31. Weber, C.; Zabinski, S.; Koschitzky, T.; et al. Transplantation 49:396-404; 1990.

Third Series of Experiments

T cell proliferation and cytokine production in diabetic NOD mice transplanted with encapsulated porcine islets.

The goal of these experiments was to develop techniques for transplanting microencapsulated xenogeneic islets as a durable physiologic source of insulin for diabetic patients. In spontaneously diabetic NOD mice, encapsulated neonatal porcine islets plus CTLA4-Ig treatment reversed diabetes for over 100 days, but encapsulated islets without CTLA4-Ig were rejected in about 2 weeks and unencapsulated islets within 1 week. (See Fig. 36)

The proliferative and cytokine responses of T cells from NOD mice transplanted with encapsulated porcine islets were compared. (see Figs. 37 through 58) Spleen cells (SPC) from rejecting NODs spontaneously proliferated in vitro, whereas SPC from mice with functioning grafts or non-transplanted NODs did not. Islet cells induced no proliferation above background with either normal or transplanted NOD SPC. However, cytokine secretion after stimulation with porcine islets was detected. (see Figs. 39A - 53B) SPC from both rejecting and non-rejecting mice secreted IFN γ , IL-10, and TGF β as well as low levels of IL-2, IL-12 and IL-4 when stimulated with islets. In addition, fluid from peritoneal cavities (the site of transplanted encapsulated islets) contained IFN γ , NO $_2$, IL-12 and high levels of TGF- β . (see Figs. 41A, 52A, and 58; 48A; and 52A,) By contrast, porcine islets stimulated no cytokine secretion by cells from control NOD mice. (see e.g. Figs. 41B, 48B, 50B, 51B)

It was a surprise to find no significant differences in islet-induced proliferation or cytokines between NOD mice that rejected or accepted grafts. Therefore, additional cytokines (for example, IL-1 and TNF α) that

05/22/00 5:38:40 PM

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

levels.

Only non-rejecting transplanted NODs have been tested for nitric oxide production. Rejecting NODs as well as normal NODs, diabetic NODs, and BALB/c mice must be analyzed for nitric oxide generation by SPC and PECs.

Lymphokines produced by spleen cells

Stimulating spleen cells from control animals (untransplanted normal NODs, diabetic NODs, or BALB/c mice) with pig islets did not induce lymphokines.

However, stimulating spleen cells from control mice with Con A induced relatively high levels of IFN- γ and IL-2, low amounts of IL-10, and no IL-4, IL-5, or TNA- α .

In over 50% of transplanted mice tested, pig islets stimulated IFN- γ secretion by spleen cells in vitro (10 of 17). Four mice (2 rejecting and 2 non-rejecting) produced relatively high IFN- γ (> 1000 pg/ml).

Pig islets stimulated lower levels (100-500 pg/ml) of IL-4, IL-10, and TGF- β . In 2 of 17, >100 pg/ml IL-4 was stimulated (both not rejecting). In 5 of 17, >100 pg/ml IL-10 was stimulated (2 not rejecting; 3 rejecting). In 2 of 17, >100 pg/ml TGF- β was stimulated (both not rejecting).

Pig islets stimulated IL-12 secretion by spleen cells from 1 of 17 transplanted mice (not rejecting). Pig islets stimulated IL-2 secretion above background levels in 1 of 17 transplanted mice (not rejecting). Pig islets did not stimulate TNF- α secretion by spleen cells from any transplanted mice.

Lymphokines that are clearly associated with graft

004985-0329

rejection have not been identified. No striking differences in lymphokines from mice given different CTLA4-Ig treatment (none, mutant, or WT) was observed.

05/07/00 09:00:00

8. The method of claim 7, wherein the viable cells are genetically engineered prior to transplantation into the subject.
9. The method of claim 1, wherein the subject is a human.
10. The method of claim 9, wherein the viable cells are derived from a mammal.
11. The method of claim 10, wherein the mammal is a human.
12. The method of claim 1, wherein the subject is a domesticated animal.
13. The method of claim 12, wherein the domesticated animal is a cow, a calf, a pig, a sheep, a lamb, a horse, or a chicken.
14. The method of claim 12, wherein the viable cells comprise cells which secrete a hormone which promotes growth in the domesticated animal.
15. The method of claim 1, wherein the viable cells comprise cells which secrete a biologically active substance.
16. The method of claim 15, wherein the cells which secrete a biologically active substance are endocrine cells.
17. The method of claim 16, wherein the endocrine cells are insulin-producing cells, hepatocytes, parathyroid cells, or pituitary cells.
18. The method of claim 15, wherein the cells which

96220"59864060

secrete a biologically active substance are neuroectodermal cells.

19. The method of claim 18, wherein the neuroectodermal cells are adrenal cells or lymphocytes.
20. The method of claim 1, wherein the semipermeable membrane is impermeable to immunoglobulins and/or lymphocytes.
21. The method of claim 2, wherein treating the subject with CTLA4 comprises administering soluble CTLA4 to the subject.
22. The method of claim 21, wherein the soluble CTLA4 is CTLA4Ig.
23. The method of claim 1, wherein inhibiting the subject's immune system from responding to said contained cells or tissue comprises inhibiting production of immunoglobulins and activated macrophages capable of reacting with the viable cells or tissue.
24. A method of treating diabetes in a subject which comprises:
 - a) containing viable insulin-producing cells, or tissue comprising viable insulin-producing cells, within a device comprising a semipermeable membrane so as to obtain contained viable insulin-producing cells;
 - b) transplanting contained viable insulin-producing cells obtained in step (a) into the subject in an amount effective to treat diabetes in the

004066 9864060

subject; and

c) treating the subject with a substance which inhibits an immune-system costimulation event in an amount effective to inhibit the subject's immune system from responding to an amount of contained viable insulin-producing cells according to step (b).

25. The method of claim 24, wherein the substance which inhibits an immune-system costimulation event is CTLA4.
26. The method of claim 24, wherein the tissue comprising the viable insulin-producing cells comprises pancreatic islet tissue.
27. The method of claim 24, wherein the viable insulin-producing cells comprise cells which have been genetically engineered prior to transplantation to secrete insulin.
28. The method of claim 24, wherein the device is a hollow fiber, a disk, or a sphere.
29. The method of claim 24, wherein the device is a microcapsule.
30. The method of claim 24, wherein the viable insulin-producing cells or the tissue comprising the viable insulin-producing cells are derived from a xenogeneic donor.
31. The method of claim 24, wherein the viable insulin-producing cells or the tissue comprising the viable insulin-producing cells are derived from an allogeneic donor.

32. The method of claim 24, wherein the viable insulin-producing cells or the tissue comprising the viable insulin-producing cells are derived from the subject.
33. The method of claim 32, wherein the viable insulin-producing cells are genetically engineered to secrete insulin prior to transplantation into the subject.
34. The method of claim 24, wherein the subject is afflicted with insulin-dependent diabetes mellitus.
35. The method of claim 34, wherein the subject is a mammal.
36. The method of claim 35, wherein the subject is a human.
37. The method of claim 24, wherein the subject is a mammal.
38. The method of claim 37, wherein the subject is a human.
39. The method of claim 24, wherein the semipermeable membrane is impermeable to immunoglobulins and/or lymphocytes.
40. The method of claim 25, wherein treating the subject with CTLA4 comprises administering soluble CTLA4 to the subject.
41. The method of claim 40, wherein the soluble CTLA4 is CTLA4Ig.

42. The method of claim 24, wherein inhibiting the subject's immune system from responding to said contained viable insulin-producing cells or tissue comprises inhibiting production of immunoglobulins and activated macrophages capable of reacting with the viable insulin-producing cells or tissue.
43. The method of claim 1, wherein the substance which inhibits an immune-system costimulation event also alters the cytokine profile of the subject so as to protect the contained cells or tissue from the subject's immune system.
44. The method of claim 43, wherein the substance increases the production of gamma-interferon in the subject.
45. The method of claim 43, wherein the substance is CTLA4Ig.
46. The method of claim 1, wherein the substance which inhibits an immune-system costimulation event binds complement.
47. The method of claim 46, wherein the substance is CTLA4Ig.
48. The method of claim 1, wherein the substance which inhibits an immune-system costimulation event does not alter the cytokine profile of the subject so as to protect the contained cells or tissue from the subject's immune system.
49. The method of claim 48, wherein the substance increases the production of gamma-interferon and IL-2 in the subject.

004966 0329
00220 59864060

50. The method of claim 48, wherein the substance is CTLA4Ig.
51. The method of claim 48, wherein the substance and the containing of the viable cells within the device comprising the semipermeable membrane prevents host immune cell proliferation in the subject.
52. The method of claim 48, wherein the device comprising the semipermeable membrane is a hollow, fiber, a disc, or a sphere.
53. The method of claim 48, wherein the device comprising the semipermeable membrane is a microcapsule.

004985 032799
062230 59867060


METHOD OF INHIBITING IMMUNE SYSTEM DESTRUCTION OF
TRANSPLANTED VIABLE CELLS

Abstract of the Disclosure

This invention provides a method of inhibiting viable cells transplanted into a subject from being destroyed by the subject's immune system which comprises: a) containing the viable cells, or tissue comprising the viable cells, prior to transplantation within a device comprising a semipermeable membrane; and b) treating the subject with a substance which inhibits an immune-system costimulation event in an amount effective to inhibit the subject's immune system from responding to said contained cells or tissue. In one embodiment, the substance which inhibits an immune-system costimulation event is CTLA4. Also provided by this invention is a method of treating diabetes in a subject which comprises: a) containing viable insulin-producing cells, or tissue comprising such cells, within a device comprising a semipermeable membrane; b) transplanting an effective amount of such contained viable insulin-producing cells into the subject; and c) treating the subject with an effective amount of a substance which inhibits an immune-system costimulation event.

09043865-032799

FIG. 2






FIG. 4




FIG. 5




FIG. 6



05/22/00 59864050


7/56

FIG. 7



004985.032798

FIG. 8






FIG. 10




FIG. 11




FIG. 12





FIG. 13




85/220 5864060

FIG. 14



85/220" 59864060

FIG. 15



17/56

FIG. 17



0004935.032798
06/22/2015 09:56:10

FIG. 18




FIG. 19




FIG. 21




FIG. 24




FIG. 25A




FIG. 25B





FIG. 25C




FIG. 25D

FIG. 26





FIG. 27






FIG. 28B





FIG. 28A

FIG. 29B



B

FIG. 29A



FIG. 30





FIG. 31



004985-032798
RECEIVED 59064060

FIG. 32



FIG. 33



FIG. 34




FIG. 35 Microencapsulated Neonatal Pig Islet Transplants into Diabetic NOD Mice
Treated with mutant CTLA4-Ig for 21 Days




FIG. 36

Neonatal Porcine Islet Xenografts in NOD Mice: Effects of CTLA-4Ig and Microencapsulation




FIG. 38
 Spontaneous Proliferation by SPC from
 NODs with rejected or functioning grafts




FIG. 39A IL-2 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 39B IL-2 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 40A IL-2 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 40B IL-2 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 41A IFN-gamma present in peritoneal fluid on sac day
Transplanted NODs




FIG. 41B IFN-gamma present in peritoneal fluid on sac day
Untransplanted mice




FIG. 42A IFN gamma secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 42B IFN-gamma secretion by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 43A IL-4 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 43B IL-4 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 44A IL-4 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 44B IL-4 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 45A IL-5 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 45B IL-5 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 46A IL-10 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 46B IL-10 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 47A IL-10 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 47B IL-10 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 48A IL-12 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 48B IL-12 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 49A IL-12 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 49B IL-12 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 50A TNF alpha present in peritoneal fluid on sac day
Transplanted NODs




FIG. 50B TNF alpha present in peritoneal fluid on sac day
Untransplanted NODs




FIG. 51A TNF- α secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 51B TNF- α secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 52A TGF beta present in peritoneal fluid on sac day
Transplanted NODs




FIG. 52B TGF beta present in peritoneal fluid on sac day
Untransplanted mice




FIG. 53A TGF- β secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 53B TGF- β secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 54A NO₂ produced by SPCs cultured with pig islets
Transplanted NODs - Not Rejecting




FIG. 54B Transplanted NODs - Rejecting




FIG. 55A

NO₂ produced by PECs after 96 hr in culture
with original encapsulated porcine islets

Transplanted NODs - Not Rejecting




FIG. 55B

Transplanted NODs - Rejecting




FIG. 56
Proliferation of SPC from NOD #1335 to pig islets or insulin
(Expt 288)




FIG. 57A Proliferation of SPC from NOD #1335
to pig islet cells in vitro (Expt 288)




FIG. 57B Proliferation of SPC from NOD #1335
Expt 288






FIG. 58 NO2 Production by Cells from NOD #1335 (Expt 288)



United States Patent & Trademark Office

Office of Initial Patent Examination – Scanning Division



Application deficiencies found during scanning:

1. Application papers are not suitable for scanning and are not in compliance with 37 CFR 1.52 because:
 - ☐ All sheets must be the same size and either A4 (21 cm x 29.7 cm) or 8-1/2" x 11". Pages _____ do not meet these requirements.
 - ☐ Papers are not flexible, strong, smooth, non-shiny, durable, and white.
 - ☐ Papers are not typewritten or mechanically printed in permanent ink on one side.
 - ☐ Papers contain improper margins. Each sheet must have a left margin of at least 2.5 cm (1") and top, bottom and right margins of at least 2.0 cm (3/4").
 - ☐ Papers contain hand lettering.
2. Drawings are not in compliance and were not scanned because:
 - ☐ The drawings or copy of drawings are not suitable for electronic reproduction.
 - ☐ All drawings sheets are not the same size. Pages must be either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
 - ☐ Each sheet must include a top and left margin of at least 2.5 cm (1"), a right margin at least 1.5 cm (9/16") and a bottom margin of at least 1.0 cm (3/8").
3. Page(s) Fig. 1, 20, 23, 24 are not of sufficient clarity, contrast and quality for electronic reproduction.
4. Page(s) _____ are missing.
5. OTHER: NO DECLARATION

502220-59061060